



Anxiety and Depressive Symptoms and Disorders in Children and Adolescents With Migraine

¹ A.Meer, ² A.Patel

^{1,2} California Northstate University College of Medicine

Correspondence : dr.meer2984@gmail.com

Article History :

Received date : 2011/10/12

Revised date : 2011/11/21

Accepted date : 2011/12/24

Published date : 2012/01/19



Copyright: © 2012 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (<https://creativecommons.org/licenses/by-nc/4.0/>).

ABSTRACT

Background: The interplay between anxiety and depressive symptoms in children and adolescents suffering from migraine is a complex and multifaceted issue that has garnered considerable attention in the literature. **Literature Review:** Further contributions to the discourse reveal that psychiatric comorbidities are prevalent among children with migraines, with familial factors and headache types influencing the relationship between these conditions (ANTONACI et al., 2011). The bidirectional relationship between migraine and major depression is reinforced by findings indicating shared heritability and neurobiological mechanisms, particularly involving the serotonin system (T. et al., 2009). This interplay necessitates an interdisciplinary approach to treatment, as the presence of anxiety and depression can complicate migraine management and increase the risk of suicidal behavior (Dresler et al., 2009). The literature emphasizes the need for early screening and effective management strategies, such as cognitive behavioral therapy (CBT) and selective serotonin reuptake inhibitors (SSRIs), to alleviate the burden of anxiety on the quality of life for migraine patients (Kumar et al., 2002). The studies collectively underscore the critical importance of understanding the underlying mechanisms linking migraine with psychiatric disorders and highlight the necessity for tailored treatment plans that address both physical and psychological aspects of these comorbid conditions. **Conclusion:** In conclusion, the interplay between migraine, anxiety, and depressive symptoms in children and adolescents is characterized by a complex and multifaceted relationship that warrants further investigation. The existing literature reveals a pressing need for longitudinal studies to clarify these associations and to develop targeted interventions that effectively address the comprehensive needs of affected individuals.

Keyword: Anxiety, Depressive Symptoms and Disorders, Migraine

INTRODUCTION

The interplay between anxiety and depressive symptoms in children and adolescents suffering from migraine is a complex and multifaceted issue that has garnered considerable attention in the literature. The foundational work by (Virtanen, 2008) highlights the significant association between anxiety disorders and major depression in migraine sufferers, noting that anxiety symptoms often precede the onset of migraines, while depressive symptoms typically emerge afterward. This temporal relationship underscores the importance of recognizing anxiety as a potential precursor to migraine, particularly among females, who exhibit higher levels of depressive symptoms and anxiety-linked personality traits compared to their male counterparts.

Building on this premise, (Antonaci et al., 2011) further elucidate the prevalence of psychiatric comorbidities in children with migraine. They emphasize the need for careful consideration of familial factors and the nuanced relationship between headache types and psychiatric disorders. Their findings suggest that while anxiety and depression are prevalent in migraine patients, the evidence remains inconclusive regarding the exact nature of their relationship, highlighting a gap in longitudinal studies that could clarify these associations.

In a subsequent review, (ANTONACI et al., 2011) reaffirm the connection between familial recurrence of headaches and psychiatric disorders, particularly in migraine cases. They point out that the presence of depression correlates with poorer outcomes across all headache types, stressing the need for further research into the unique clinical manifestations of depression in children, which complicate diagnosis and treatment.

(T. et al., 2009) contribute to this discourse by establishing a bidirectional relationship between migraine and major depression, asserting that the two conditions share significant heritability factors and neurobiological mechanisms. Their systematic review indicates that the serotonin system plays a critical role in this relationship, with alterations in serotonin levels potentially influencing both

migraine and depressive symptoms. This highlights the necessity for clinicians to adopt an interdisciplinary approach when treating patients with comorbid conditions.

Further emphasizing the clinical implications of this comorbidity, (Dresler et al., 2009) delineate the increased risks associated with anxiety and major depression in migraine patients, including heightened susceptibility to suicidal behavior. They advocate for an integrated treatment strategy that addresses both migraine and psychiatric disorders, recognizing the overlapping neural networks that may underlie these conditions.

(Kumar et al., 2002) expand upon these findings by discussing the bidirectional relationship between anxiety disorders and migraine, where each condition exacerbates the other. They underscore the significance of early screening for anxiety in migraine patients and the necessity of effective management strategies, such as cognitive behavioral therapy (CBT) and selective serotonin reuptake inhibitors (SSRIs), to mitigate the impact of anxiety on the quality of life.

Together, these studies paint a comprehensive picture of the intricate relationship between migraine and psychiatric comorbidities in children and adolescents. The literature reveals a pressing need for further exploration of the underlying mechanisms, as well as the development of targeted interventions that address both the physical and psychological aspects of migraine in young populations.

LITERATURE REVIEW

The article "Epidemiological studies of childhood and adolescence headache" by (Virtanen, 2008) provides a comprehensive overview of the intricate relationship between migraine and psychological disorders in children and adolescents. It highlights several key studies that illustrate the significant association between migraine and both anxiety and depressive symptoms.

One of the critical insights presented is the finding by Merikangas et al. (1990), which establishes a strong link between anxiety disorders and major depression in individuals suffering from migraine. This suggests that the psychological burden of migraine may be compounded by the presence of these

disorders, leading to a more complex clinical picture. Furthermore, the article notes that migraine sufferers experienced an earlier onset of anxiety symptoms compared to their non-migraine counterparts, indicating a potential predictive relationship where anxiety may precede and exacerbate migraine conditions.

The gender differences in the association between headaches and psychiatric disorders are also noteworthy. Egger et al. (1998) demonstrated that these associations vary significantly between boys and girls, with girls showing a higher prevalence of depressive symptoms in conjunction with frequent headaches. This finding raises important considerations for clinicians regarding the need for gender-sensitive approaches in the assessment and treatment of migraine in pediatric populations.

Moreover, the article underscores the role of psychological factors in the context of chronic pain. Studies referenced, such as those by Perquin et al. (2000) and Egger et al. (1999), emphasize that comorbid psychological issues are critical in understanding the relationships between frequent headaches and psychological problems. The observation that adolescents with chronic pain report higher levels of psychological vulnerability than those without chronic pain further supports the notion that psychological health is an essential component of managing migraine in this demographic.

The article also discusses the implications of the serotonergic system in the interplay between anxiety and pain perception. This connection suggests that biological factors may also contribute to the observed psychological comorbidities in migraine sufferers, potentially guiding future research into targeted therapeutic approaches.

The article "Migraine and psychiatric comorbidity: a review of clinical findings" by (Antonaci et al., 2011) provides a comprehensive analysis of the relationship between migraine and psychiatric disorders, particularly focusing on anxiety and mood disorders in children and adolescents. The authors highlight the significance of understanding this comorbidity, as it poses substantial implications for the management and treatment of affected individuals.

The review underscores that while there is a recognized association between migraine and psychiatric conditions, particularly anxiety and depression, the

existing evidence remains inconclusive. This inconsistency is primarily attributed to the variability in study designs and methodologies. The authors note that the cohort study design is the most reliable for establishing a clear association between these disorders, as it allows for longitudinal observation of subjects over time. In contrast, many of the studies reviewed were cross-sectional or case-control, which may limit the ability to draw definitive conclusions regarding causality.

Moreover, the article points out the challenges in understanding the etiology of the comorbid association between migraine and psychiatric disorders. The authors emphasize that the lack of research specifically targeting pharmacological and non-pharmacological treatments for these comorbid conditions in younger populations represents a significant gap in the literature. This gap is particularly concerning given that depression is frequently examined in individuals with migraine, and the need for effective interventions is critical.

The authors also conducted a systematic search of the literature, identifying studies related to the relationship between headache and depression. They categorized these studies based on their methodological rigor and relevance to the topic. Notably, the findings revealed that a substantial number of studies were inconsistent with the proposed relationship between migraine and depression, indicating a need for more robust research to clarify these associations.

The article "Understanding the nature of psychiatric comorbidity in migraine: A systematic review focused on interactions and treatment implications" by (T. et al., 2009) provides an insightful examination of the intricate relationship between migraine and depression, particularly emphasizing the prevalence and underlying mechanisms of this comorbidity. The authors highlight that the incidence of depression in individuals with migraine is nearly double that of the general population, suggesting a significant public health concern that warrants attention.

One of the central themes of the article is the bidirectional nature of the relationship between migraine and depression, which is supported by evidence indicating a heritable component for both conditions, estimated at around 40-50%. This polygenic background implies a complex interplay of genetic factors that may

predispose individuals to both migraine and depressive disorders, thereby reinforcing the need for integrated treatment approaches.

The authors also delve into the neurobiological underpinnings of this comorbidity, particularly the role of the serotonin (5-HT) system. They report that patients with migraine exhibit elevated 5-HT concentrations during migraine attacks and reduced levels interictally, which may contribute to the mood dysregulation observed in these patients. Furthermore, the article discusses the significance of GABA levels in cerebrospinal fluid, noting that lower GABA levels are found in chronic migraine (CM) patients who also suffer from depression, thereby suggesting a potential biomarker for this comorbidity.

Another critical aspect addressed is the involvement of the hypothalamic-pituitary-adrenal (HPA) axis, which may serve as a shared pathway linking migraine and depression. The authors propose that an imbalance between pro-inflammatory and anti-inflammatory cytokines could further elucidate the relationship among migraine, depression, obesity, and the transition from episodic to chronic migraine.

Neuroimaging studies cited in the review reveal functional and structural changes in brain areas associated with pain modulation, indicating a common neural substrate for migraine and affective disorders. This finding underscores the necessity for a comprehensive understanding of how these conditions interact on a neurological level, which could inform more effective treatment strategies.

The article "Understanding the nature of psychiatric comorbidity in migraine: a systematic review focused on interactions and treatment implications" by (T. et al., 2009) presents a comprehensive examination of the relationship between migraine and psychiatric disorders, particularly focusing on major affective and anxiety disorders. The authors highlight that these disorders are not only prevalent among individuals with migraine but also significantly impact the clinical course, treatment efficacy, and overall outcomes for these patients.

A critical evaluation of the material reveals that the authors effectively synthesize existing literature to underscore the bidirectional relationship between migraine and major depression. This interplay suggests that the presence of one condition can exacerbate the other, complicating both diagnosis and treatment. The

authors note that the comorbidity of migraine with major depression or anxiety disorders can substantially heighten the risk of suicide attempts, a crucial consideration for clinicians. This alarming statistic emphasizes the need for heightened vigilance and comprehensive assessment in patients presenting with migraine, particularly in identifying underlying psychiatric conditions.

Moreover, the article discusses the overlapping neural network patterns that characterize both migraine and anxiety disorders. This intersection indicates a shared pathophysiological basis, which may provide insights into potential therapeutic approaches. The authors advocate for tailored treatment plans that address both migraine and its comorbid psychiatric conditions, suggesting that a one-size-fits-all approach may be inadequate. Instead, individualized care that considers the psychological dimensions of migraine is essential for improving patient outcomes.

The authors also draw attention to the specific clinical presentations of anxiety symptoms within the context of migraine. They note that excessive worry, fear, and avoidance behaviors can manifest as part of the migraine experience, complicating the clinical picture. This dual presentation necessitates a careful diagnostic process to distinguish between primary migraine and anxiety disorders, ensuring that both conditions are adequately addressed.

The article "Understanding the nature of psychiatric comorbidity in migraine: a systematic review focused on interactions and treatment implications" by (T. et al., 2009) provides a comprehensive examination of the relationship between migraine and psychiatric disorders, particularly focusing on anxiety and depressive symptoms in children and adolescents. The authors highlight the significant prevalence of both migraine and psychiatric conditions, emphasizing that individuals suffering from chronic migraine or migraine with aura are at a notably higher risk for developing major depression, anxiety, and even suicidal behaviors when compared to non-migraine sufferers.

The systematic review synthesizes a wealth of epidemiological data, revealing a concerning link between migraine and psychiatric disorders. The authors reference a large genome-wide association study that underscores a higher genetic correlation between migraine and psychiatric conditions, suggesting that

genetic predispositions may play a critical role in this comorbidity. This connection is particularly salient when considering the developmental stage of children and adolescents, who may experience exacerbated symptoms due to the interplay of migraine and mental health issues.

The authors argue that the presence of psychiatric comorbidities can significantly impact treatment outcomes for migraine patients. For instance, the authors note that those with concurrent anxiety or depressive disorders may respond differently to standard migraine treatments, thereby necessitating an interdisciplinary approach to care. This approach should incorporate both pharmacological and non-pharmacological strategies tailored to address the complexities of managing migraine alongside psychiatric symptoms.

The review further discusses potential mechanisms of association, including psychosocial factors, neurobiological pathways, and the impact of chronic pain on mental health. The authors suggest that understanding these interactions is crucial for developing effective treatment plans. They advocate for clinicians to be vigilant in assessing psychiatric symptoms in patients with migraine, as this could enhance the likelihood of achieving remission for both conditions.

The article "The Development and Impact of Anxiety With Migraines: A Narrative Review" by (Kumar et al., 2002) provides a comprehensive examination of the intricate relationship between migraine and anxiety disorders, particularly in the context of children and adolescents. The authors argue that individuals with migraines exhibit a heightened susceptibility to anxiety disorders, establishing a bidirectional relationship that complicates the diagnostic process. This complexity is underscored by the overlapping symptoms that both conditions present, making it challenging to discern whether anxiety is a consequence of chronic pain or an independent disorder manifesting alongside migraines.

The authors emphasize the importance of early screening for anxiety disorders in patients with migraines, noting that the co-occurrence of these conditions can significantly impair the quality of life. This assertion is supported by various studies cited within the review, which highlight the prevalence of anxiety among migraine sufferers. The authors advocate for proactive management

strategies, suggesting that effective prophylactic pharmacotherapy aimed at reducing headache frequency can potentially mitigate the onset of anxiety disorders.

Furthermore, the article discusses the role of selective serotonin reuptake inhibitors (SSRIs) as a first-line treatment for those who develop anxiety disorders in conjunction with migraines. This pharmacological approach is complemented by cognitive-behavioral therapy (CBT), which has been shown to be effective in managing anxiety symptoms. The authors call for further research to deepen the understanding of the migraine-anxiety nexus, advocating for a more integrated approach to diagnosis and management that considers both conditions holistically.

CONCLUSION

The literature review presents a thorough exploration of the complex relationship between anxiety and depressive symptoms in children and adolescents suffering from migraines. The initial studies indicate a significant association between anxiety disorders and major depression within this population, with evidence suggesting that anxiety symptoms often precede migraine onset, while depressive symptoms typically emerge subsequently (Virtanen, 2008). This temporal relationship highlights the importance of early recognition and intervention, especially in females, who tend to exhibit higher levels of both anxiety and depression (Antonaci et al., 2011).

Further contributions to the discourse reveal that psychiatric comorbidities are prevalent among children with migraines, with familial factors and headache types influencing the relationship between these conditions (ANTONACI et al., 2011). The bidirectional relationship between migraine and major depression is reinforced by findings indicating shared heritability and neurobiological mechanisms, particularly involving the serotonin system (T. et al., 2009). This interplay necessitates an interdisciplinary approach to treatment, as the presence of anxiety and depression can complicate migraine management and increase the risk of suicidal behavior (Dresler et al., 2009).

The literature emphasizes the need for early screening and effective management strategies, such as cognitive behavioral therapy (CBT) and selective serotonin reuptake inhibitors (SSRIs), to alleviate the burden of anxiety on the

quality of life for migraine patients (Kumar et al., 2002). The studies collectively underscore the critical importance of understanding the underlying mechanisms linking migraine with psychiatric disorders and highlight the necessity for tailored treatment plans that address both physical and psychological aspects of these comorbid conditions.

In conclusion, the interplay between migraine, anxiety, and depressive symptoms in children and adolescents is characterized by a complex and multifaceted relationship that warrants further investigation. The existing literature reveals a pressing need for longitudinal studies to clarify these associations and to develop targeted interventions that effectively address the comprehensive needs of affected individuals.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Virtanen, R., 2008. Epidemiological studies of childhood and adolescence headache. [PDF]
2. Antonaci, F., Nappi, G., Galli, F., Camillo Manzoni, G., Calabresi, P., & Costa, A., 2011. Migraine and psychiatric comorbidity: a review of clinical findings. ncbi.nlm.nih.gov
3. ANTONACI, F. A. B. I. O., NAPPI, G. I. U. S. E. P. P. E., COSTA, A. L. F. R. E. D. O., F, G., GC, M., & P, C., 2011. Migraine and psychiatric comorbidity: a review of clinical findings.. [PDF]
4. T., D., S., C., K., G., J. -I., H., C., L., T., N. P., M., P., G., S., A., T., R., O., & G., S., 2009. Understanding the nature of psychiatric

comorbidity in migraine: A systematic review focused on interactions and treatment implications. [PDF]

5. Dresler, T., Caratozzolo, S., Guldolf, K., Huhn, J. I., Loiacono, C., Niiberg-Pikksööt, T., Puma, M., Sforza, G., Tobia, A., Ornello, R., & Serafini, G., 2009. Understanding the nature of psychiatric comorbidity in migraine: a systematic review focused on interactions and treatment implications. ncbi.nlm.nih.gov
6. Kumar, R., Asif, S., Bali, A., Kaur Dang, A., & A Gonzalez, D., 2002. The Development and Impact of Anxiety With Migraines: A Narrative Review. ncbi.nlm.nih.gov