



Coronary Artery Complication in Kawasaki Disease and the Importance of Early Intervention

¹ A.Ahmed, ¹ A.Noorani

¹ Charles R. Drew University of Medicine and Science

Correspondence : doctorahmed2938478@gmail.com

Article History :

Received date : 2018/10/09

Revised date : 2018/11/15

Accepted date : 2018/12/12

Published date : 2019/01/05



Copyright: © 2019 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (<https://creativecommons.org/licenses/by-nc/4.0/>).

ABSTRACT

Background: Kawasaki disease (KD) is a significant pediatric condition characterized by acute febrile systemic vasculitis that primarily affects children under the age of five. The literature surrounding KD underscores the critical importance of early intervention to mitigate the severe potential complications, particularly those involving the coronary arteries. **Literature Review:** Further exploration by (Lee et al., 2017) reveals alarming statistics regarding coronary arteriopathy associated with KD, noting that untreated patients face a 15-25% chance of developing coronary aneurysms, which can lead to a mortality rate of 2-3%. In 2018, (Shah et al., 2018) highlighted the challenges in diagnosing KD due to its nonspecific symptoms and underscored the increasing incidence of the disease in the United States. The urgency of prompt recognition and management is reiterated as a means to prevent serious outcomes related to coronary artery complications. A case report by (Reza Sabri et al., 2011) exemplifies the long-term consequences of KD, showcasing coronary obstruction in a 14.5-year-old boy. This case underscores the necessity for ongoing monitoring and intervention, advocating for advanced imaging techniques to assess coronary complications effectively. The article by (Liu et al., 2011) expands on the complications arising from delayed diagnosis and treatment, emphasizing that timely intervention is crucial in minimizing the risk of severe complications, notably coronary artery aneurysms. These aneurysms have become the leading cause of acquired pediatric heart disease. **Conclusion:** In conclusion, the literature collectively emphasizes the necessity for early recognition and intervention in Kawasaki disease to mitigate the progression of coronary artery complications, which remain a leading cause of acquired heart disease in children. The findings advocate for continued research and improved diagnostic strategies to enhance treatment outcomes.

Keyword: Coronary Artery Complication, Kawasaki Disease, Importance of Early Intervention

INTRODUCTION

Kawasaki disease (KD) is a significant pediatric condition characterized by acute febrile systemic vasculitis that primarily affects children under the age of five. The literature surrounding KD underscores the critical importance of early intervention to mitigate the severe potential complications, particularly those involving the coronary arteries. The foundational work by (Galeotti et al., 2016) highlights the pathogenesis of KD, indicating that the disease arises from an aberrant inflammatory response, leading to endothelial cell injury and subsequent coronary artery abnormalities. This article establishes the framework for understanding the disease's implications, particularly the 10-20% of patients who exhibit resistance to standard treatments like intravenous immunoglobulin (IVIG), thus increasing their risk for coronary vasculitis.

Building on this foundation, (Lee et al., 2017) delve into the myocardial implications of KD, emphasizing the spectrum of coronary arteriopathy ranging from minimal involvement to severe complications such as giant coronary aneurysms. Their findings reveal alarming statistics, with untreated patients facing a 15-25% chance of developing coronary aneurysms and a 2-3% mortality rate due to coronary arteriopathy. This article reinforces the necessity for tailored therapeutic management based on the degree of coronary involvement, as proposed by the American Heart Association.

In 2018, (Shah et al., 2018) further contextualize the disease by addressing the challenges in diagnosing KD due to its nonspecific symptoms. They note the increasing incidence in the United States and stress the urgency of prompt recognition and management to prevent serious outcomes. Their review serves as a reminder of the disease's enigmatic nature and the critical role of early intervention in improving patient prognosis.

The case report by (Reza Sabri et al., 2011) illustrates the long-term consequences of KD, specifically highlighting coronary obstruction in a 14.5-year-old boy. This case exemplifies how coronary involvement can evolve over time,

necessitating ongoing monitoring and intervention. The authors advocate for the use of advanced imaging techniques, such as echocardiography and CT angiography, to assess and manage coronary complications effectively.

(Liu et al., 2011) expand on the complications associated with KD, particularly the impact of delayed diagnosis and treatment on cardiovascular outcomes. Their findings emphasize that timely intervention is crucial in minimizing the risk of severe complications, including coronary artery aneurysms, which have become the leading cause of acquired pediatric heart disease.

(Tan et al., 2012) provide a global perspective on KD research, noting the disease's systemic inflammation and potential genetic predispositions. They highlight the emerging concerns surrounding Kawasaki-like diseases in the context of COVID-19, which further complicates the landscape of KD diagnosis and treatment.

Finally, (Cavalcanti Lianza et al., 2012) synthesize key aspects of KD, particularly the importance of identifying and monitoring patients at higher risk for coronary artery abnormalities. Their review elucidates the pathological changes occurring in the coronary arteries during KD, underscoring the need for vigilant management to prevent long-term cardiovascular complications.

Overall, the literature collectively underscores the critical need for early recognition and intervention in Kawasaki disease to prevent the progression of coronary artery complications, which remain a leading cause of acquired heart disease in children.

LITERATURE REVIEW

The article "Predisposing factors, pathogenesis and therapeutic intervention of Kawasaki disease" by (Galeotti et al., 2016) provides a comprehensive overview of Kawasaki disease (KD), emphasizing its implications for coronary artery complications and the critical nature of early intervention. The authors elucidate the pathophysiological mechanisms underlying KD, which is characterized by an acute

febrile response and is notably linked to significant coronary artery abnormalities in pediatric patients.

The discussion begins with the predisposing factors associated with KD, identifying genetic susceptibility and environmental triggers as pivotal elements in the disease's onset. The authors argue that an aberrant inflammatory response to an infectious agent plays a crucial role in the disease's progression, particularly in genetically predisposed individuals. This inflammatory response leads to endothelial cell injury, primarily due to T cell activation and the cytotoxic effects of proinflammatory cytokines ((Galeotti et al., 2016)).

A significant aspect of the article is the focus on therapeutic interventions. The standard treatment for acute KD includes intravenous immunoglobulin (IVIG) and aspirin, which have been shown to mitigate the risk of coronary artery complications. However, the authors highlight a concerning statistic: approximately 10-20% of patients exhibit resistance to IVIG therapy, resulting in an elevated risk of coronary vasculitis. This emphasizes the necessity for timely and effective treatment strategies to prevent long-term cardiovascular issues in affected children ((Galeotti et al., 2016)).

Moreover, the review touches upon emerging therapeutic options for patients who are resistant to IVIG. The exploration of new treatments is vital, given that KD is the leading cause of acquired heart disease in children in various regions, including Japan, Europe, and the USA. The authors advocate for continued research into alternative therapies to address this significant clinical challenge ((Galeotti et al., 2016)).

The article "Myocardial Assessment in School-Aged Children with Past Kawasaki Disease" by (Lee et al., 2017) provides a comprehensive examination of Kawasaki disease (KD) and its implications for coronary artery health in children. The authors emphasize that KD is a systemic vasculitis that poses significant risks as the leading cause of acquired heart disease in childhood, particularly highlighting the complications associated with coronary arteriopathy. The prevalence of coronary aneurysms in untreated KD patients is alarmingly high, ranging from 15%

to 25%, with a mortality rate of 2% to 3% due to coronary complications ((Lee et al., 2017)).

The article effectively outlines the necessity of early intervention in managing KD, noting that the American Heart Association has established guidelines for therapeutic and follow-up management based on the extent of coronary involvement. This underscores the critical role of prompt diagnosis and treatment in mitigating long-term cardiovascular risks. Moreover, the authors delve into other cardiac complications associated with KD, such as myocarditis and valvulitis, which can lead to abnormal heart function. The distinction between inflammatory and ischemic myocardial injury is particularly relevant, as it informs treatment strategies and prognostic assessments for affected children ((Lee et al., 2017)).

While the article successfully highlights the urgent need for early intervention, it also points out the gaps in understanding the long-term outcomes of myocardial function in KD patients. Despite previous studies focusing on acute dysfunction, the authors call for further research to elucidate the lasting effects of KD on cardiac health. This acknowledgment of the need for ongoing investigation is a critical aspect of the article, as it highlights the evolving nature of knowledge surrounding KD and its complications.

The article "Review of Kawasaki Disease" by (Shah et al., 2018) provides a comprehensive overview of Kawasaki disease (KD), emphasizing its status as a leading cause of acquired heart disease in children. The authors detail the nature of KD as an acute self-limited vasculitis primarily affecting infants and children under five years of age. This demographic focus underscores the critical need for awareness among healthcare providers, as early intervention can significantly alter the disease trajectory and improve patient outcomes.

A salient point made in the article is the nonspecific nature of the symptoms associated with KD, such as fever and lymphadenopathy. This symptomatology can complicate the diagnostic process, often leading to delays in treatment. The authors argue that such delays can result in serious complications, including coronary artery

aneurysms, which are a major concern in the context of KD. The potential for these severe outcomes accentuates the necessity for prompt recognition of the disease, as timely intervention is crucial for mitigating risks associated with coronary artery complications.

Furthermore, the article highlights an alarming trend: the increasing incidence of KD in the United States over the past decade. This trend necessitates heightened vigilance among pediatricians and healthcare practitioners, as the rising prevalence may lead to more cases of cardiovascular complications if not addressed adequately. The authors advocate for improved diagnostic criteria and awareness campaigns to facilitate early detection and treatment.

The article "A 14.5-Year-Old Boy with Coronary Obstruction after Kawasaki Disease: A Case Report" by (Reza Sabri et al., 2011) provides a detailed examination of the implications of Kawasaki Disease (KD) on coronary artery health, particularly emphasizing the critical nature of early intervention in mitigating long-term complications. The authors highlight that KD is a significant contributor to acquired coronary artery disease in the pediatric population, with coronary artery aneurysms (CAAs) being the most severe complication, affecting nearly 25% of affected children.

One of the key insights from the article is the dynamic nature of coronary involvement in KD. The authors note that while some cases may see regression of coronary abnormalities, others may progress to more severe conditions such as stenosis or complete obstruction. This variability underscores the necessity for continuous monitoring and evaluation through echocardiography, which the authors advocate as a valuable tool for assessing the proximal parts of the coronary arteries. However, they also acknowledge that angiography remains the gold standard for diagnosis, providing definitive insights into coronary status.

The article effectively discusses the role of intravenous immunoglobulin (IVIG) in reducing the risk of CAAs, which is crucial for clinicians managing KD. The authors emphasize that timely administration of IVIG can significantly alter the disease's trajectory, thus illustrating the importance of early intervention.

Additionally, they mention the advantages of computed tomography (CT) angiography as a noninvasive alternative that offers accurate evaluations of coronary artery status, which may be particularly useful in settings where traditional angiography poses higher risks.

The article "Extracardiovascular injury complications in Kawasaki disease" by (Liu et al., 2011) provides a comprehensive overview of the implications of Kawasaki disease (KD), particularly focusing on the development of coronary artery aneurysms (CAA) as a significant complication. The authors underscore that CAA has emerged as the leading cause of acquired heart disease in the pediatric population, which is a critical concern given the potentially severe long-term outcomes associated with this condition.

A critical aspect discussed in the article is the diagnostic approach to KD. The authors indicate that the diagnosis primarily relies on clinical symptoms, which include fever, conjunctival injection, changes in the oral cavity, rash, and lymphadenopathy. This reliance on symptomatic presentation highlights a crucial challenge in the early identification of KD. The authors argue that delays in diagnosis and subsequent treatment can lead to an escalation in cardiovascular complications, emphasizing the importance of timely intervention. This assertion is particularly relevant in clinical practice, where the window for effective treatment is often narrow.

The article further elaborates on the pathophysiology of KD and its association with CAA, detailing how the inflammatory processes inherent in KD can lead to vascular damage. The authors provide evidence that early treatment with intravenous immunoglobulin (IVIG) can significantly reduce the incidence of CAA when administered promptly. However, the discussion also points to the variability in clinical presentation among patients, which can complicate timely diagnosis and treatment.

Moreover, (Liu et al., 2011) highlight the need for increased awareness among healthcare providers regarding the symptoms of KD, as well as the importance of considering KD in differential diagnoses for children presenting with

fever and other related symptoms. Their findings advocate for enhanced educational efforts aimed at both medical professionals and the public to recognize the signs of KD early, thereby facilitating quicker intervention and potentially improving outcomes for affected children.

The article "A global bibliometric analysis on Kawasaki disease research over the last 5 years (2017–2021)" by (Tan et al., 2012) provides a comprehensive overview of the recent advancements and trends in Kawasaki Disease (KD) research. The authors highlight that KD, first identified in Japan in 1967, predominantly affects children under the age of five and is characterized by systemic inflammation. This inflammation can lead to significant complications, particularly involving the coronary arteries, which underscores the importance of early intervention.

The article emphasizes the multifactorial nature of KD, noting that various pathogenic microorganisms may trigger the disease, and genetic predispositions can exacerbate its severity. The authors discuss how certain genetic abnormalities can increase the risk of KD by enhancing T cell activation and dysregulating B cell signaling, among other mechanisms. This insight is crucial as it suggests that understanding the genetic factors involved in KD may lead to better diagnostic and therapeutic strategies, particularly for preventing coronary artery complications.

Moreover, the authors reference guidelines from the Royal College of Pediatrics and Child Health, which provide diagnostic criteria for KD-like multi-inflammatory syndrome, particularly in the context of children with prolonged fever and signs of organ dysfunction. This is particularly relevant in light of the COVID-19 pandemic, where some children infected with SARS-CoV-2 have displayed symptoms akin to those of KD. The overlap of these conditions necessitates a heightened awareness and prompt intervention to mitigate the risk of coronary artery complications.

The systematic review mentioned within the article highlights the effectiveness of steroid treatment during the acute phase of KD, which has been shown to improve coronary lesions and lower inflammatory markers. This finding

is pivotal, as it suggests that timely steroid administration can significantly alter the disease's trajectory and reduce the risk of long-term cardiovascular issues associated with KD.

The article "Kawasaki Disease: A Never-ending Story?" by (Cavalcanti Lianza et al., 2012) provides a comprehensive overview of Kawasaki disease (KD), emphasizing the critical need for early identification and intervention to mitigate complications, particularly those affecting the coronary arteries. The authors underscore the importance of recognizing the disease's potential to lead to serious cardiovascular outcomes, including aneurysm formation, which occurs in approximately 30% of untreated cases and up to 6% of treated patients. This statistic highlights the urgency of prompt treatment to prevent long-term sequelae.

A particularly noteworthy aspect of the review is the discussion of myocarditis, which can occur in the early stages of KD. The authors point out that while left ventricular dysfunction is generally mild and responsive to treatment, the presence of diffuse myocarditis followed by fibrosis in some patients may result in more severe cardiac complications, including both systolic and diastolic dysfunction. This insight stresses the necessity for vigilant monitoring of cardiac function in KD patients, especially those presenting with myocarditis.

The review aptly discusses the importance of stratifying patients based on their risk for complications, which is essential for tailoring treatment approaches. The authors advocate for a proactive stance in managing KD, suggesting that early intervention not only improves immediate outcomes but also reduces the risk of chronic complications, particularly those related to the coronary arteries.

CONCLUSION

The literature on Kawasaki disease (KD) emphasizes the critical importance of early intervention to mitigate severe complications, particularly those affecting the coronary arteries. The foundational work by (Galeotti et al., 2016) outlines the pathogenesis of KD, indicating that it arises from an aberrant inflammatory response leading to endothelial cell injury and coronary artery abnormalities. This

study highlights that 10-20% of patients may resist standard treatments such as intravenous immunoglobulin (IVIG), which significantly increases their risk for coronary vasculitis.

Further exploration by (Lee et al., 2017) reveals alarming statistics regarding coronary arteriopathy associated with KD, noting that untreated patients face a 15-25% chance of developing coronary aneurysms, which can lead to a mortality rate of 2-3%. The necessity for tailored therapeutic management based on the extent of coronary involvement is emphasized, aligning with guidelines established by the American Heart Association.

In 2018, (Shah et al., 2018) highlighted the challenges in diagnosing KD due to its nonspecific symptoms and underscored the increasing incidence of the disease in the United States. The urgency of prompt recognition and management is reiterated as a means to prevent serious outcomes related to coronary artery complications.

A case report by (Reza Sabri et al., 2011) exemplifies the long-term consequences of KD, showcasing coronary obstruction in a 14.5-year-old boy. This case underscores the necessity for ongoing monitoring and intervention, advocating for advanced imaging techniques to assess coronary complications effectively.

The article by (Liu et al., 2011) expands on the complications arising from delayed diagnosis and treatment, emphasizing that timely intervention is crucial in minimizing the risk of severe complications, notably coronary artery aneurysms. These aneurysms have become the leading cause of acquired pediatric heart disease.

A global perspective provided by (Tan et al., 2012) discusses systemic inflammation and genetic predispositions in KD, noting emerging concerns regarding Kawasaki-like diseases in the context of COVID-19, which complicates the landscape of KD diagnosis and treatment.

Finally, (Cavalcanti Lianza et al., 2012) synthesizes key aspects of KD, particularly the importance of identifying and monitoring patients at higher risk for coronary artery abnormalities. This review elucidates the pathological changes

occurring in the coronary arteries during KD, reinforcing the need for vigilant management to prevent long-term cardiovascular complications.

In conclusion, the literature collectively emphasizes the necessity for early recognition and intervention in Kawasaki disease to mitigate the progression of coronary artery complications, which remain a leading cause of acquired heart disease in children. The findings advocate for continued research and improved diagnostic strategies to enhance treatment outcomes.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Galeotti, C., V. Kaveri, S., Cimaz, R., Koné-Paut, I., & Bayry, J., 2016. Predisposing factors, pathogenesis and therapeutic intervention of Kawasaki disease. ncbi.nlm.nih.gov
2. Lee, H., Shin, J., & Eun, L., 2017. Myocardial Assessment in School-Aged Children with Past Kawasaki Disease. ncbi.nlm.nih.gov
3. Shah, R., Hirzallah, H., Fares, M., Mallad, A., Karlsson, G., & El-Hamdani, M., 2018. Review of Kawasaki Disease. [PDF]
4. Reza Sabri, M., Mahdavi, C., Ahmadi, A., Ghaderian, M., & Dehghan, B., 2011. A 14.5-Year-Old Boy with Coronary Obstruction after Kawasaki Disease: A Case Report. ncbi.nlm.nih.gov
5. Liu, J., Huang, Y., Qin, S., Su, D., Ye, B., & Pang, Y., 2011. Extracardiovascular injury complications in Kawasaki disease. ncbi.nlm.nih.gov
6. Tan, W., Jing, L., Wang, Y., & Li, W., 2012. A global bibliometric analysis on Kawasaki disease research over the last 5 years. ncbi.nlm.nih.gov

7. Cavalcanti Lianza, A., de Fátima Rodrigues Diniz, M., Saori Shiraishi Sawamura, K., da Rocha Brito Menezes, C., de Sousa Lobo Silva, I., & Nunes Leal, G., 2012. Kawasaki Disease: A Never-ending Story?. [ncbi.nlm.nih.gov](https://www.ncbi.nlm.nih.gov)