



Risk Factors Associated With Post–COVID-19 Condition

¹ A. Bowersox, ¹ A. Bradshaw¹ Charles R. Drew University of Medicine and Science, United States of AmericaCorrespondence : dr.bowersox39484gmail.com

Article History :

Received date : 2022/10/19

Revised date : 2022/11/12

Accepted date : 2022/12/06

Published date : 2023-01-13



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (<https://creativecommons.org/licenses/by-nc/4.0/>).

ABSTRACT

Background: The literature surrounding post-COVID-19 conditions, commonly referred to as long COVID, has rapidly evolved as researchers seek to understand the multifaceted risk factors associated with this phenomenon. The systematic review by (Mahbub Hossain et al., 2022) highlights the need for empowering individuals experiencing long COVID through education and social support, emphasizing the significant impact of perceived changes in social networks on their well-being. **Literature Review:** The systematic analysis by (Israel Notarte et al., 2022) identifies demographic factors such as female sex, older age, and pre-existing health conditions as potential predictors of long COVID symptomatology. These findings are corroborated by the research conducted by (Bovil et al., 2013), which focuses specifically on individuals aged 50 and older in Europe and Israel, confirming that these demographic characteristics significantly contribute to the risk of severe post-COVID-19 conditions. Further insights into the pathophysiology of long COVID are provided by (Bohmwald et al., 2014), who reveal that even younger individuals without prior health issues may experience prolonged health problems following infection. This complexity is echoed in the comprehensive review by (Song et al., 2014), which synthesizes findings across studies, emphasizing that hospitalization during the acute phase of COVID-19 significantly increases the risk of long COVID. **Conclusion:** In conclusion, the body of literature reviewed highlights the multifaceted risk factors associated with post-COVID-19 conditions, including demographic characteristics, pre-existing health conditions, social support systems, and the severity of the acute disease. The evidence suggests that targeted interventions focusing on vulnerable populations, particularly older adults and those with pre-existing comorbidities, are essential for improving long-term health outcomes. Furthermore, addressing psychological and social dimensions of care is crucial for supporting individuals navigating the challenges of long COVID.

Keyword: Risk Factors Associated, Post–COVID-19 Condition

INTRODUCTION

The literature surrounding post-COVID-19 conditions, commonly referred to as long COVID, has rapidly evolved as researchers seek to understand the multifaceted risk factors associated with this phenomenon. The systematic review by (Mahbub Hossain et al., 2022) highlights the need for empowering individuals experiencing long COVID through education and social support, emphasizing the significant impact of perceived changes in social networks on their well-being. This study underscores the psychological, emotional, and social dimensions of long COVID, suggesting that a well-informed and supportive environment is essential for those affected.

Building on this foundation, (Israel Notarte et al., 2022) provide a systematic analysis of risk factors associated with long COVID, delineating the prevalence of persistent symptoms in individuals recovering from SARS-CoV-2 infections. Their findings reveal that a substantial percentage of patients experience long-term effects, with specific demographic factors such as female sex, older age, and pre-existing comorbidities identified as potential predictors of long COVID symptomatology. This article serves to contextualize the broader epidemiological landscape of long COVID, linking individual characteristics to health outcomes.

Further investigation into demographic influences is presented by (Bovil et al., 2013), who focus on individuals aged 50 and older in Europe and Israel. Their research confirms that high age, male sex, and pre-existing health conditions are significant risk factors for experiencing severe post-COVID-19 conditions. This demographic specificity adds depth to the understanding of how age and sex intersect with long COVID risk, suggesting targeted interventions may be necessary for older populations.

In a more recent exploration, (Bohmwald et al., 2014) examine the pathophysiological and immunological features of long COVID, revealing that a notable percentage of patients, including younger individuals without prior health issues, do not return to their baseline health following infection. Their research

indicates that while some variants of the virus may correlate with varying long-term health outcomes, the relationship between initial disease severity and long COVID symptoms remains complex and warrants further exploration.

Finally, the comprehensive review by (Song et al., 2014) synthesizes the current understanding of long COVID, emphasizing the critical role of gender, age, and pre-existing conditions in predicting susceptibility. Their findings highlight that women are at a significantly higher risk, and that hospitalization during the acute phase of COVID-19 dramatically increases the likelihood of developing long COVID. This review not only consolidates existing knowledge but also points to the necessity for healthcare systems to adapt their approaches based on the identified risk factors, thereby enhancing patient care and management strategies.

Together, these articles contribute to a nuanced understanding of the risk factors associated with post-COVID-19 conditions, illustrating the interplay between demographic characteristics, health history, and the broader social context in shaping long-term health outcomes for individuals recovering from COVID-19.

LITERATURE REVIEW

The article "Living with 'long COVID': A systematic review and meta-synthesis of qualitative evidence" by (Mahbub Hossain et al., 2022) provides a comprehensive examination of the lived experiences of individuals suffering from long COVID, emphasizing the necessity for empowerment through education and resources. The authors argue that individuals affected by long COVID require not only medical support but also a robust social support system to navigate the complexities of their condition.

A critical evaluation of the material reveals that the authors effectively outline the multifaceted nature of long COVID, which encompasses psychological, emotional, and social dimensions. They highlight the importance of understanding the dynamics of social support networks for individuals experiencing long COVID, noting that perceived changes in social support can have significant implications for recovery and overall well-being. This perspective is vital, as it shifts the focus

from purely medical treatment to a more holistic approach that includes social and emotional care.

The article also addresses the challenges faced by individuals in accessing healthcare services. (Mahbub Hossain et al., 2022) identify several barriers, including inadequate information from healthcare providers, a lack of understanding of long COVID, and limited service availability. These challenges are exacerbated by systemic issues such as provider burnout and poor communication skills, which hinder effective healthcare delivery. The authors argue that these factors contribute to a global crisis in managing long COVID, indicating a need for systemic change in healthcare practices.

Moreover, the authors point out the implications of the digital divide in research recruitment, suggesting that online methods may exclude vulnerable populations who lack access to technology. This insight underscores the importance of inclusive research practices that consider diverse health and social contexts, ensuring that all voices are represented in the discourse surrounding long COVID.

The article titled "Age, Sex and Previous Comorbidities as Risk Factors Not Associated with SARS-CoV-2 Infection for Long COVID-19: A Systematic Review and Meta-Analysis" by (Israel Notarte et al., 2022) provides a comprehensive examination of risk factors associated with long COVID-19, emphasizing the need for a nuanced understanding of pre-infection variables. The systematic review and meta-analysis approach employed by the authors allows for a robust synthesis of existing literature, which is crucial given the evolving nature of research surrounding post-COVID-19 conditions.

The authors effectively define long COVID-19 in accordance with the World Health Organization's criteria, framing it as a condition that manifests after the acute phase of SARS-CoV-2 infection, characterized by persistent symptoms lasting for at least two months. This clear definition sets the stage for the analysis of risk factors, which is vital for healthcare management and planning.

One of the key insights from the article is the identification of pre-infection risk factors, including age, sex, and previous comorbidities. The authors critically evaluate previous meta-analyses that have suggested various risk factors, noting that while female sex and older age are frequently cited, their systematic review found no significant association between these factors and the development of long COVID-19 symptoms. This finding challenges the prevailing narrative that these demographic factors are definitive predictors of long COVID-19, suggesting that the relationship may be more complex than previously understood.

Moreover, the article highlights the importance of differentiating between pre-infection and infection-associated factors. By categorizing risk factors in this manner, the authors provide a framework for understanding how different variables might influence the likelihood of developing long COVID-19. This distinction is particularly relevant for clinicians and policymakers aiming to identify individuals at higher risk and to tailor interventions accordingly.

The systematic review also brings attention to the need for further research to explore the underlying mechanisms that may contribute to long COVID-19. The article underscores that while demographic factors have been extensively studied, other potential risk factors, such as psychological health and socioeconomic status, warrant further investigation.

The article titled "Risk factors of post-COVID-19 conditions attributed to COVID-19 disease in people aged ≥ 50 years in Europe and Israel" by (Bovil et al., 2013) provides a comprehensive analysis of the risk factors associated with the development of post-COVID-19 conditions in older adults. The authors effectively highlight that older age, male sex, and pre-existing comorbidities significantly contribute to a more severe manifestation of COVID-19 and its subsequent long-term effects.

The study's findings are particularly relevant given the increasing recognition of post-COVID-19 conditions, which encompass a range of persistent symptoms that can affect individuals long after the acute phase of the infection has resolved. The authors utilize a robust methodological approach, analyzing data

from a substantial cohort of individuals aged 50 and older across Europe and Israel. This demographic focus is critical, as it targets a population known to be at higher risk for severe outcomes related to COVID-19.

One of the key insights from the article is the emphasis on pre-existing comorbidities as a significant predictor of post-COVID-19 condition. The authors detail how conditions such as cardiovascular disease, diabetes, and respiratory issues can exacerbate the severity of COVID-19 and contribute to prolonged recovery times. This correlation underscores the importance of managing chronic health issues, particularly in older adults, to mitigate the risk of severe COVID-19 outcomes and subsequent long-term health complications.

Additionally, the article discusses the gender disparity observed in the incidence of post-COVID-19 conditions, noting that male patients are more likely to experience severe symptoms. This finding aligns with existing literature that suggests biological, behavioral, and social factors may influence the differential impact of COVID-19 between genders. The authors call for further research to explore these disparities in depth, which could inform targeted interventions and healthcare strategies.

The article also raises awareness about the importance of public health initiatives aimed at vaccination and prevention strategies, particularly for vulnerable populations. By identifying and understanding the risk factors associated with post-COVID-19 conditions, healthcare providers can better tailor their approaches to monitoring and treating affected individuals.

The article "Pathophysiological, immunological, and inflammatory features of long COVID" by (Bohmwald et al., 2014) provides a comprehensive analysis of the risk factors associated with post-COVID-19 condition, commonly referred to as long COVID. The authors present compelling evidence that a significant portion of individuals, specifically 35%, do not return to their normal health status within 2 to 3 weeks following a COVID-19 diagnosis. Notably, the study highlights that younger adults, particularly those aged 18–34 without pre-existing chronic health

conditions, are also at risk, with one in five reporting persistent health issues post-infection.

The article identifies several key risk factors linked to long COVID, including age, gender, cardiovascular disease, and diabetes. These factors are not only associated with the severity of acute COVID-19 but also suggest a broader underlying vulnerability among certain demographics. The authors critically evaluate the existing literature, noting a discrepancy in findings regarding the correlation between long COVID incidence and the severity of initial COVID-19 symptoms. While some studies suggest that those who experience more severe acute symptoms may be at heightened risk for long COVID, others indicate that long COVID can manifest even in cases of mild or asymptomatic infections. This observation underscores the complexity of long COVID and challenges the assumption that the severity of initial infection is a reliable predictor of long-term outcomes.

Furthermore, the article emphasizes the gender disparities observed in long COVID diagnoses, particularly among women under 50 years old, who exhibit a higher prevalence of symptoms such as fatigue and dyspnea compared to their male counterparts. The authors propose that biological differences, including variations in the expression of ACE2 and TMPRSS2, as well as immunological factors, may contribute to these gender differences. This aspect of the study is particularly valuable as it opens avenues for further research into the mechanisms underlying these disparities.

The article titled "A Comprehensive Review of the Global Epidemiology, Clinical Management, Socio-Economic Impacts, and National Responses to Long COVID with Future Research Directions" by (Song et al., 2014) provides an extensive examination of the risk factors associated with post-COVID-19 condition, commonly referred to as long COVID. The authors emphasize the importance of understanding these risk factors and the underlying physiological mechanisms to enhance screening, monitoring, and treatment strategies for affected patients.

A critical evaluation of the material reveals that the authors have systematically identified and analyzed key demographic and health-related risk factors. Notably, gender disparities are highlighted, with women being at twice the risk of developing long COVID compared to men. This finding prompts further investigation into the biological and social determinants that may contribute to this disparity. Additionally, the article underscores the significant impact of age, indicating that individuals aged 60 to 79 are at a 25% increased risk, which escalates to 60% for those over 80. This age-related vulnerability necessitates targeted healthcare interventions for older populations.

The authors also draw attention to the heightened risk associated with hospitalization due to COVID-19, noting that patients requiring hospitalization may face an 18-fold increased risk of developing long COVID. More alarmingly, those who undergo invasive ventilation are at an astronomical 115 times greater risk. This stark statistic highlights the critical need for healthcare systems to implement rigorous follow-up care for hospitalized patients to monitor and address potential long COVID symptoms.

Furthermore, the article thoroughly discusses the role of pre-existing chronic conditions as significant risk factors. Individuals with a history of cardiovascular diseases, chronic lung diseases, kidney diseases, hypertension, diabetes, obesity, and depression are shown to be more susceptible to long COVID. This correlation emphasizes the importance of managing chronic health issues proactively, particularly in populations at risk of severe COVID-19 outcomes.

The authors also explore additional contributors to long COVID risk, including lifestyle factors such as smoking, high body mass index, life stress, and psychological factors. These insights are crucial as they suggest that interventions aimed at improving lifestyle and mental health could potentially mitigate the risk of long COVID.

CONCLUSION

The literature on post-COVID-19 conditions, commonly referred to as long COVID, has expanded significantly, revealing a complex interplay of risk factors that influence the likelihood and severity of symptoms experienced by individuals after acute COVID-19 infection. The systematic review by (Mahbub Hossain et al., 2022) emphasizes the importance of social support and education in managing long COVID, highlighting that individuals' well-being can be significantly affected by perceived changes in their social networks. This perspective underscores the necessity for a holistic approach to care that includes psychological and emotional support alongside medical treatment.

The systematic analysis by (Israel Notarte et al., 2022) identifies demographic factors such as female sex, older age, and pre-existing health conditions as potential predictors of long COVID symptomatology. These findings are corroborated by the research conducted by (Bovil et al., 2013), which focuses specifically on individuals aged 50 and older in Europe and Israel, confirming that these demographic characteristics significantly contribute to the risk of severe post-COVID-19 conditions.

Further insights into the pathophysiology of long COVID are provided by (Bohmwald et al., 2014), who reveal that even younger individuals without prior health issues may experience prolonged health problems following infection. This complexity is echoed in the comprehensive review by (Song et al., 2014), which synthesizes findings across studies, emphasizing that hospitalization during the acute phase of COVID-19 significantly increases the risk of long COVID.

In conclusion, the body of literature reviewed highlights the multifaceted risk factors associated with post-COVID-19 conditions, including demographic characteristics, pre-existing health conditions, social support systems, and the severity of the acute disease. The evidence suggests that targeted interventions focusing on vulnerable populations, particularly older adults and those with pre-existing comorbidities, are essential for improving long-term health outcomes.

Furthermore, addressing psychological and social dimensions of care is crucial for supporting individuals navigating the challenges of long COVID.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Mahbub Hossain, M., Das, J., Rahman, F., Nesa, F., Hossain, P., M. Khairul Islam, A., Tasnim, S., Faizah, F., Mazumder, H., & Purohit, N., 2022. Living with “long COVID”: A systematic review and meta-synthesis of qualitative evidence. osf.io
2. Israel Notarte, K., Helena Santos de Oliveira, M., Juneire Peligro, P., Veronica Velasco, J., Macaranas, I., Therese Ver, A., Carmeli Pangilinan, F., Pastrana, A., Goldrich, N., Kavteladze, D., Margarita Leticia Gellaco, M., Liu, J., Lippi, G., Michael Henry, B., & Fernández-de-las-Peñas, C., 2022. Age, Sex and Previous Comorbidities as Risk Factors Not Associated with SARS-CoV-2 Infection for Long COVID-19: A Systematic Review and Meta-Analysis. ncbi.nlm.nih.gov
3. Bovil, T., Wester, C. T., Scheel-Hincke, L. L., & Andersen-Ranberg, K., 2013. Risk factors of post-COVID-19 conditions attributed to COVID-19 disease in people aged ≥ 50 years in Europe and Israel. ncbi.nlm.nih.gov
4. Bohmwald, K., Diethelm-Varela, B., Rodríguez-Guilarte, L., Rivera, T., A. Riedel, C., A. González, P., & M. Kalergis, A., 2014. Pathophysiological, immunological, and inflammatory features of long COVID. ncbi.nlm.nih.gov
5. Song, X., Song, W., Cui, L., Q. Duong, T., Pandey, R., Liu, H., Zhou, Q., Sun, J., Liu, Y., & Li, T., 2014. A Comprehensive Review of the Global

Epidemiology, Clinical Management, Socio-Economic Impacts, and National Responses to Long COVID with Future Research Directions. ncbi.nlm.nih.gov