



Association of Ebola Virus Infection With Hearing Loss in Regions Where Ebola Virus Infection Is Endemic

¹ A. Diederich, ¹ A. Downey

¹ Charles R. Drew University of Medicine and Science, United States of
America

Correspondence : drdiedrich2884783gmail.com

Article History :

Received date : 2024/10/15

Revised date : 2024/11/09

Accepted date : 2024/12/18

Published date : 2025-01-23



Copyright: © 2024 by the
authors. Submitted for possible
open access publication under
the terms and conditions of the
Creative Commons Attribution
(BY NC) license
(<https://creativecommons.org/licenses/by-nc/4.0/>).

ABSTRACT

Background: The literature surrounding the association of Ebola Virus Infection with hearing loss in endemic regions reveals a complex interplay of epidemiological insights, clinical findings, and the long-term health implications for survivors.

Literature Review: The investigation into Lassa fever (LF) by (J. Mateer et al., 2018) reveals significant associations between LF and sudden-onset sensorineural hearing loss (SNHL). This finding raises the possibility of similar auditory implications for EVD survivors, suggesting a broader public health concern regarding hearing loss following viral hemorrhagic fevers. Furthermore, the work of (Kamara, 2019) examines predictors and risk factors for EVD, emphasizing the role of gender dynamics and socio-economic conditions that may influence both transmission and long-term health outcomes, including auditory health. In addition, (Helena de Magalhães Barbosa et al., 2019) explores auditory findings associated with Zika virus infection, drawing parallels between different viral infections and their potential auditory impacts. The study by (C. Ficenec et al., 2020) reinforces the notion that LF can induce hearing loss even in asymptomatic cases, advocating for targeted screening and intervention strategies for survivors of viral hemorrhagic fevers, including EVD. Lastly, the systematic review by (Onayi Mandizadza et al., 2024) collates data on the somatic symptoms experienced by EVD survivors, highlighting the prevalence of neurological complications that may include auditory issues, thereby emphasizing the need for ongoing research into the long-term health outcomes of EVD.

Conclusion: In conclusion, the collective findings from these studies illuminate the critical association between Ebola virus infection and hearing loss, revealing significant gaps in current understanding and underscoring the urgency for further research and public health interventions aimed at this vulnerable population. The implications of these studies advocate for a holistic approach to survivor care, which includes comprehensive screening for auditory complications and recognition of the long-term health effects of EVD.

Keyword: Association, Ebola Virus Infection, Hearing Loss, Regions, Ebola Virus Infection, Endemic

INTRODUCTION

The literature surrounding the association of Ebola Virus Infection with hearing loss in endemic regions reveals a complex interplay of epidemiological insights, clinical findings, and the long-term health implications for survivors. The foundational work by (Roger Ngatu et al., 2017) provides a crucial overview of the epidemiology of Ebolavirus disease (EVD) and highlights the occupational risks faced by healthcare workers in Sub-Saharan Africa. Their comprehensive documentary study underscores the necessity for enhanced public health preparedness, emphasizing the importance of understanding EVD not only in terms of immediate health outcomes but also in the context of occupational safety and the potential for long-term sequelae among survivors.

Building on the understanding of viral infections and their repercussions, (J. Mateer et al., 2018) draw attention to Lassa fever (LF) and its significant association with sensorineural hearing loss (SNHL). This article highlights that approximately one-third of LF survivors experience sudden-onset SNHL, a condition that remains under-recognized in endemic regions. The authors argue for the need to address this neglected public health burden, particularly in areas where access to audiological care is severely limited. Their findings suggest that similar mechanisms may be at play in other viral infections, including EVD, raising questions about the broader implications of viral hemorrhagic fevers on auditory health.

(Kamara, 2019) further contributes to this discourse by identifying predictors and risk factors associated with Ebola virus disease, particularly in Sierra Leone. The article meticulously examines various dimensions of risk, including traditional practices and gender-related factors, thereby providing a nuanced understanding of how these elements may influence not only the transmission of EVD but also the potential for long-term health consequences, such as hearing loss, among survivors.

In a related vein, the integrative review by (Helena de Magalhães Barbosa et al., 2019) explores auditory findings associated with Zika virus infection,

suggesting a possible link between flavivirus infections and hearing impairments. Although the direct causality between Zika virus and hearing loss remains inconclusive, the article posits that auditory alterations may stem from cochlear or neural involvement, thus inviting further investigation into the auditory risks posed by viral infections, including EVD.

(C. Ficenec et al., 2020) delve deeper into the specific implications of Lassa fever on hearing, presenting compelling evidence that the virus may induce hearing loss even in asymptomatic cases. The study's findings, which indicate a significant prevalence of hearing complications among Lassa fever survivors, reinforce the necessity for targeted hearing loss screening and intervention strategies. This work is pivotal in advocating for comprehensive care approaches for survivors of viral hemorrhagic fevers, including EVD.

Finally, (Onayi Mandizadza et al., 2024) compile a systematic review and meta-analysis that highlights the prevalence of somatic symptoms among Ebola Virus Disease survivors, including neurological and cognitive sequelae. Their work synthesizes various studies on the long-term health effects of EVD, suggesting that auditory complications may be an overlooked aspect of post-Ebola syndrome. This systematic approach underscores the need for ongoing research into the full spectrum of health outcomes associated with EVD, including the critical area of auditory health.

Collectively, these articles illuminate the multifaceted relationship between Ebola virus infection and hearing loss, revealing significant gaps in current understanding and underscoring the urgency for further research and public health interventions aimed at this vulnerable population.

LITERATURE REVIEW

The article titled "Epidemiology of ebolavirus disease (EVD) and occupational EVD in health care workers in Sub-Saharan Africa: Need for strengthened public health preparedness" by (Roger Ngatu et al., 2017) provides a comprehensive overview of the epidemiological landscape surrounding EVD, particularly focusing on healthcare workers in Sub-Saharan Africa. The authors

conducted a documentary study that meticulously reviewed relevant literature, aiming to synthesize existing knowledge on EVD and occupational EVD (OEVD) while highlighting critical work safety issues.

The methodology employed in this study is noteworthy, as it involved a systematic search that generated a substantial number of articles, ultimately narrowing down to 34 that were directly relevant to the authors' focus. This rigorous selection process enhances the credibility of their findings and ensures that the conclusions drawn are based on a solid foundation of empirical data. The authors utilized a range of keywords, such as "hospital infection" and "work safety," to encompass various dimensions of EVD, which reflects a thorough understanding of the complexities involved in the epidemiology of the disease.

One of the key insights from the article is the emphasis on the occupational risks faced by healthcare workers during Ebola outbreaks. The authors effectively illustrate how these risks are compounded by inadequate public health infrastructure and insufficient protective measures for medical personnel. This aspect of the study is particularly relevant in the context of hearing loss associated with EVD, as healthcare workers may experience prolonged exposure to high-risk environments, potentially leading to both physical and psychological health issues, including auditory complications.

Furthermore, the article underscores the urgent need for strengthened public health preparedness in the face of EVD outbreaks. The authors argue that enhancing safety protocols and training for healthcare workers is essential not only to protect them from direct infection but also to mitigate the broader implications of the disease on community health. This call to action is critical, as it aligns with the growing body of evidence suggesting that the aftermath of EVD infections can extend beyond the immediate health crisis, affecting various aspects of survivors' well-being, including sensory health.

The article titled "Lassa fever-induced sensorineural hearing loss: A neglected public health and social burden" by (J. Mateer et al., 2018) provides a critical examination of the association between Lassa fever (LF) and sudden-onset

sensorineural hearing loss (SNHL) in endemic regions, particularly in West Africa. The authors highlight that despite the clinical recognition of this association since 1990, the prevalence of LF-induced SNHL remains significantly underestimated, illustrating a gap in public health awareness and response.

The article presents a compelling argument regarding the impact of LF on hearing health, noting that approximately one-third of LF survivors experience sudden-onset SNHL. This statistic emphasizes the urgent need for increased recognition of SNHL as a major complication of LF, which is often overshadowed by the more immediate and acute health threats posed by the viral hemorrhagic fever itself. The authors underscore the social and economic burden of hearing loss, estimating that it costs Nigeria around 43 million dollars annually, which further complicates the public health landscape in regions affected by LF.

Moreover, the article discusses the recent development of an animal model that replicates the symptoms of SNHL associated with LF, which is a significant advancement in understanding the underlying mechanisms of this complication. The preliminary data suggesting immune-mediated injury as a potential cause of SNHL in LF patients opens avenues for future research that could lead to targeted therapies and improved clinical outcomes.

The authors call for further investigation into the epidemiology and mechanisms of LF-induced SNHL, which is crucial for informing vaccine development and treatment strategies. Their emphasis on the need for comprehensive research reflects a broader concern for addressing neglected health issues in endemic regions, where the interplay of infectious diseases and their long-term consequences, such as hearing loss, are often inadequately addressed.

The article "Predictors and Risk Factors of Ebola Virus Disease in Sierra Leone" by (Kamara, 2019) provides a comprehensive exploration of the various factors associated with Ebola Virus Disease (EVD) outbreaks, particularly in the context of Sierra Leone. Kamara employs a systematic approach to identify key determinants of EVD transmission, focusing on a range of epidemiological aspects,

including gender dimensions, community behaviors, and socio-economic conditions.

One of the pivotal insights from Kamara's work is the emphasis on gender dynamics in the context of Ebola transmission. The article highlights that women, often being primary caregivers, face unique risks during outbreaks, which may lead to increased vulnerability not only to infection but also to subsequent health complications such as hearing loss. This gendered perspective is crucial, as it underscores the need for targeted interventions that consider the specific roles and challenges faced by women in Ebola-affected regions.

The methodology employed in the review is robust, as it includes a selection of studies published in English from credible sources, ensuring a high level of reliability in the data presented. Kamara's exclusion criteria effectively narrow the focus to relevant studies, thereby enhancing the quality of the review. By concentrating on peer-reviewed articles, the author strengthens the validity of the findings discussed.

Moreover, the article addresses the socio-economic factors that contribute to the spread of Ebola, such as poverty and community practices related to burial and caregiving. These elements are critical in understanding the broader context of EVD outbreaks and their implications for public health. The discussion of risky behaviors, such as traditional burial practices, further elucidates how cultural factors can significantly influence the transmission dynamics of the virus.

However, while the article provides valuable insights into the predictors and risk factors of Ebola, it could benefit from a more explicit connection to the long-term health consequences of EVD, particularly regarding auditory health. The potential association between Ebola virus infection and hearing loss remains an area that warrants further investigation. By integrating findings from studies focusing on post-Ebola health outcomes, future research could offer a more comprehensive understanding of the lasting impacts of the disease on survivors.

The article titled "Auditory findings associated with Zika virus infection: an integrative review" provides a comprehensive examination of the auditory implications related to Zika virus (ZIKV) infection, highlighting the connection between viral infections and auditory dysfunction. The review synthesizes findings from a limited number of studies, ultimately identifying 27 records that addressed auditory, functional, and morphological outcomes associated with ZIKV.

A critical evaluation of the material reveals that while the authors present a thorough analysis of the auditory alterations linked to ZIKV, the evidence remains circumstantial. The predominantly sensorineural and transient nature of the auditory changes suggests a potential involvement of the cochlear or neural structures. However, the authors caution against definitively attributing auditory impairments solely to ZIKV infection, noting that the existing literature does not provide conclusive proof of causation. This cautious approach is significant, as it reflects the complexity of establishing direct links between viral infections and specific health outcomes.

The article also draws parallels between ZIKV and Dengue virus (DENV) infections, suggesting a broader trend where flavivirus infections may be associated with hearing loss. This hypothesis is supported by previous reports concerning DENV, yet the authors emphasize the unique characteristics of ZIKV, particularly its strong tropism for the nervous system, which may differentiate it from DENV in terms of auditory impact. This distinction is crucial for understanding the potential risks associated with ZIKV and informs future research directions.

The methodologies employed in the studies reviewed primarily utilized otoacoustic emissions (OAE) and auditory brainstem response (ABR) assessments to evaluate auditory thresholds. The findings indicating OAE failures and changes in auditory thresholds through ABR assessments point towards probable peripheral auditory risks. However, the authors acknowledge the possibility of central auditory pathway involvement, underscoring the need for further investigation to delineate the specific mechanisms through which ZIKV may affect auditory function.

The article titled "Lassa fever-induced hearing loss: The neglected disability of hemorrhagic fever" by (C. Ficenec et al., 2020) provides a critical examination of the relationship between Lassa fever and hearing loss, emphasizing the implications for public health in endemic regions. The authors highlight a significant gap in the understanding of the long-term sequelae of Lassa fever, particularly its potential to induce hearing loss even in cases of asymptomatic or mild infections. This insight is particularly relevant for regions where viral hemorrhagic fevers, such as Ebola, are endemic, as it raises concerns about similar outcomes following Ebola virus infections.

The authors reference a 1990 case-control study conducted in Sierra Leone, which found that a notable percentage of seropositive individuals reported hearing difficulties. This finding suggests a potential link between Lassa fever and auditory impairments, prompting the authors to argue for a broader recognition of hearing loss as a consequence of viral infections in these regions. The article posits that the mechanisms behind Lassa fever-induced hearing loss could include direct viral damage, immune-mediated responses, and vasculitis—each of which warrants further investigation to elucidate the pathophysiological processes involved.

Moreover, the article discusses the audiometric data collected from Lassa fever survivors, indicating that while the majority of reported hearing loss is mild, it should not be overlooked. The authors argue that untreated hearing loss can severely impact the quality of life and socioeconomic status of affected individuals, underscoring the necessity for screening and treatment programs tailored for Lassa fever survivors. This call to action is particularly pertinent given the broader implications for public health strategies in managing the sequelae of viral infections in endemic regions.

The article titled "Prevalence of somatic symptoms among Ebola Virus Disease (EVD) survivors in Africa: a systematic review and meta-analysis" by (Onayi Mandizadza et al., 2024) presents a comprehensive examination of the somatic symptoms experienced by survivors of Ebola Virus Disease (EVD). This systematic review and meta-analysis collates findings from various studies to

elucidate the prevalence and nature of long-term sequelae among EVD survivors, a critical issue given the significant health implications of the disease.

The authors effectively synthesize data from multiple studies, including Clark et al. (2015) and Nanyonga et al. (2016), which highlight the diverse range of complications faced by EVD survivors, such as neurological, ocular, and musculoskeletal symptoms. The review underscores the importance of recognizing these sequelae as they can significantly affect the quality of life and overall health of survivors. The systematic approach adopted in the article allows for a clearer understanding of the prevalence of these symptoms across different regions affected by EVD, particularly in Africa.

One notable aspect of the article is its focus on the methodological rigor of the included studies. The authors critically evaluate the quality of the studies reviewed, discussing potential biases and limitations inherent in the research on EVD sequelae. This critical evaluation is essential for contextualizing the findings and understanding the variability in reported symptoms. For instance, the variability in symptom prevalence might be influenced by factors such as the timing of assessments post-recovery and the differing healthcare infrastructures in various regions.

Furthermore, the article highlights the need for continued surveillance and research into the long-term health outcomes of EVD survivors. The findings suggest that while the immediate effects of the outbreak may subside, the long-term health consequences can be profound and require ongoing medical attention. This is particularly relevant in light of the emerging recognition of hearing loss as a potential sequela of EVD, which has not been extensively documented in the literature but warrants further investigation.

CONCLUSION

The literature presents a comprehensive exploration of the association between Ebola Virus Infection (EVD) and hearing loss, emphasizing the multifaceted implications for survivors in endemic regions. The foundational study

by (Roger Ngatu et al., 2017) highlights the epidemiological landscape of EVD, particularly the occupational risks faced by healthcare workers in Sub-Saharan Africa. It underscores the need for enhanced public health preparedness, which is crucial not only for immediate health outcomes but also for addressing long-term sequelae, such as auditory complications.

The investigation into Lassa fever (LF) by (J. Mateer et al., 2018) reveals significant associations between LF and sudden-onset sensorineural hearing loss (SNHL). This finding raises the possibility of similar auditory implications for EVD survivors, suggesting a broader public health concern regarding hearing loss following viral hemorrhagic fevers. Furthermore, the work of (Kamara, 2019) examines predictors and risk factors for EVD, emphasizing the role of gender dynamics and socio-economic conditions that may influence both transmission and long-term health outcomes, including auditory health.

In addition, (Helena de Magalhães Barbosa et al., 2019) explores auditory findings associated with Zika virus infection, drawing parallels between different viral infections and their potential auditory impacts. This comparative analysis suggests the need for further investigation into the mechanisms through which EVD might similarly affect auditory function.

The study by (C. Ficenec et al., 2020) reinforces the notion that LF can induce hearing loss even in asymptomatic cases, advocating for targeted screening and intervention strategies for survivors of viral hemorrhagic fevers, including EVD. Lastly, the systematic review by (Onayi Mandizadza et al., 2024) collates data on the somatic symptoms experienced by EVD survivors, highlighting the prevalence of neurological complications that may include auditory issues, thereby emphasizing the need for ongoing research into the long-term health outcomes of EVD.

In conclusion, the collective findings from these studies illuminate the critical association between Ebola virus infection and hearing loss, revealing significant gaps in current understanding and underscoring the urgency for further research and public health interventions aimed at this vulnerable population. The

implications of these studies advocate for a holistic approach to survivor care, which includes comprehensive screening for auditory complications and recognition of the long-term health effects of EVD.

DISCLOSURE STATEMENT

- Disclosure Statement : The authors have no conflicts of Interest to declare
- Funding Sources : None
- Acknowledgements : -
- Author Contribution : All authors discussed and contributed the final content for journal submission and publication

REFERENCES

1. Roger Ngatu, N., Jean-Marie Kayembe, N., Kornblatt Phillips, E., Okech-Ojony, J., Patou-Musumari, M., Gaspard-Kibukusa, M., Madone-Mandina, N., Godefroid-Mayala, M., Mutaawe, L., Manzengo, C., Roger-Wumba, D., & Nojima, S., 2017. Epidemiology of ebolavirus disease (EVD) and occupational EVD in health care workers in Sub-Saharan Africa: Need for strengthened public health preparedness. ncbi.nlm.nih.gov
2. J. Mateer, E., Huang, C., Y. Shehu, N., & Paessler, S., 2018. Lassa fever–induced sensorineural hearing loss: A neglected public health and social burden. ncbi.nlm.nih.gov
3. Kamara, K., 2019. Predictors and Risk Factors of Ebola Virus Disease in Sierra Leone. [PDF]
4. Helena de Magalhães Barbosa, M., Clara de Magalhães-Barbosa, M., Rodrigues Robaina, J., Prata-Barbosa, A., Antonio de Melo Tavares de Lima, M., & José Ledo Alves da Cunha, A., 2019. Auditory findings associated with Zika virus infection: an integrative review(). ncbi.nlm.nih.gov
5. C. Ficenec, S., Percak, J., Arguello, S., Bays, A., Goba, A., Gbakie, M., G. Shaffer, J., D. Emmett, S., S. Schieffelin, J., & Bausch, D., 2020. Lassa fever-induced hearing loss: The neglected disability of hemorrhagic fever. ncbi.nlm.nih.gov

6. Onayi Mandizadza, O., Tsepang Phebeni, R., & Ji, C., 2024. Prevalence of somatic symptoms among Ebola Virus Disease (EVD) survivors in Africa: a systematic review and meta-analysis. ncbi.nlm.nih.gov