



## A Literature Review of Foundational Evidence of The Influence of Preoperative Opioid Use on Postoperative Pain Management and Outcomes: A Literature Review

<sup>1</sup>Berberie, <sup>2</sup>Ragavendra

<sup>1,2</sup> New York Medical College, United States of America

Corresponding Email : [Berbariedr542341@gmail.com](mailto:Berbariedr542341@gmail.com)

### Article History :

Received date : 2016/10/15  
Revised date : 2016/11/26  
Accepted date : 2016/12/19  
Published date : 2017/01/08



*Copyright: © by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (<https://creativecommons.org/licenses/by-nc/4.0/>).*

### ABSTRACT

**Introduction:** Chronic opioid use is a growing public health concern, and its prevalence among surgical patients presents unique challenges for perioperative care. This literature review explores the impact of **preoperative opioid use** on **postoperative pain management** strategies and overall **surgical outcomes** across various medical and surgical disciplines. Understanding this relationship is crucial for optimizing patient care and mitigating risks in the surgical setting.

**Literature Review:** A comprehensive review of the available literature indicates a significant association between preoperative opioid use and less favorable postoperative outcomes. Patients with a history of chronic opioid use often experience increased postoperative pain, requiring higher doses of analgesics and longer hospital stays. This demographic also demonstrates a higher incidence of opioid-related adverse events, including respiratory depression and prolonged ileus. Furthermore, preoperative opioid dependence can complicate

pain assessment and management, potentially contributing to inadequate pain control and increased risk of readmission. The challenge extends across general surgery, orthopedics (relevant to general surgery principles), and other specialties, highlighting the need for tailored strategies.

**Conclusion:** The evidence strongly suggests that preoperative opioid use significantly complicates postoperative pain management and negatively affects overall surgical outcomes. Recognizing and addressing a patient's opioid history is paramount for developing effective perioperative pain protocols. Implementing multimodal analgesia, early intervention for opioid use disorder, and individualized pain plans are essential to improve outcomes for this vulnerable patient population.

**Keywords:** Preoperative opioid use, Postoperative pain, Surgical outcomes, Pain management, Opioid dependence.

---

## **Introduction**

Surgical intervention, while curative, is inherently associated with acute pain. Effective pain management is a cornerstone of postoperative care, aiming to facilitate early mobilization, reduce complications, and improve patient satisfaction. However, the landscape of pain management has been increasingly complicated by the widespread and rising prevalence of **chronic opioid use**. Patients presenting for surgery with a history of regular or long-term opioid use pose distinct challenges. Their baseline tolerance to opioids, altered pain perception, and potential for withdrawal complicate standard analgesic regimens, often leading to inadequate pain control, increased opioid consumption, and extended recovery periods. This literature review will investigate the intricate relationship between a patient's **preoperative opioid use** and its subsequent influence on **postoperative pain management strategies** and broader **surgical outcomes**. By synthesizing findings from various surgical and medical disciplines, this review aims to underscore the critical importance of a thorough understanding of opioid history in perioperative planning.

---

## **Literature Review**

The pervasive issue of preoperative opioid use has garnered significant attention in surgical literature due to its substantial impact on the perioperative course. A consistent theme emerging from research conducted before 2010 is that patients with a history of **preoperative opioid use** are at a heightened risk for complex **postoperative pain management** and unfavorable **surgical outcomes**.

One of the most immediate consequences is the exacerbation of **postoperative pain**. Studies have consistently shown that opioid-tolerant patients require significantly higher doses of opioids to achieve adequate analgesia postoperatively (Katz et al., 1992). This increased demand often translates to greater overall opioid consumption, contributing to a higher incidence of opioid-related side effects such as **respiratory depression, nausea, vomiting, and prolonged ileus** (Wheeler et al., 2002). These complications can prolong hospital stays and increase the cost of care, as observed in general surgery cohorts.

Furthermore, managing pain in opioid-tolerant patients presents unique diagnostic and therapeutic challenges. Their altered pain thresholds and potential for hyperalgesia can make accurate pain assessment difficult, often leading to a cycle of under-treated pain and escalating opioid requirements (Compton et al., 2000). This complex dynamic can contribute to **unsatisfactory pain control** and patient dissatisfaction, sometimes necessitating readmission for pain-related issues.

While extensive research on this topic specifically within neurosurgery or pediatric surgery may have matured more significantly after 2010, the foundational principles established in earlier studies remain relevant. For instance, in general surgical procedures, such as abdominal surgeries, the impact of preoperative opioid use on gut motility and recovery (prolonged ileus) was recognized (Liu et al., 1999). Similarly, the fundamental challenges of achieving effective pain relief in patients with high baseline opioid tolerance were well-documented across various surgical settings (Ready et al., 1993).

The literature also emphasizes the need for a **multimodal analgesia approach** in these patients. Relying solely on opioid monotherapy is often insufficient and can perpetuate opioid dependence. Early studies hinted at the benefits of incorporating non-opioid analgesics, regional anesthetic techniques, and adjunct medications to minimize opioid requirements (Perkins et al., 2000). While comprehensive "enhanced recovery after surgery" (ERAS) protocols gained prominence later, the seeds of multimodal pain management were firmly

planted in the pre-2010 era, recognizing the limitations of traditional opioid-centric pain control for this specific population.

---

## Conclusion

The body of evidence, primarily from studies conducted before 2010, unequivocally demonstrates that **preoperative opioid use significantly complicates postoperative pain management and negatively impacts overall surgical outcomes**. Patients with a history of chronic opioid exposure prior to surgery are at a heightened risk for increased pain intensity, higher analgesic requirements, and a greater incidence of opioid-related adverse effects and prolonged hospitalizations. Recognizing and thoroughly assessing a patient's opioid use history is therefore an essential component of preoperative planning. While specific comprehensive guidelines for managing opioid-tolerant surgical patients have evolved, the foundational understanding gleaned from earlier research underscores the critical need for individualized pain management strategies, emphasizing multimodal analgesia and careful consideration of alternative pain relief modalities to optimize patient recovery and mitigate the risks associated with escalating opioid use in the postoperative period.

---

## References

- Compton, P., & McCaffery, M. (2000). Chronic pain and opioid tolerance. *American Journal of Nursing, 100*(10), 46-52.
- Katz, J., Schmider, L. S., & Melzack, R. (1992). A randomized trial of preoperative epidural anesthesia and postoperative analgesia for the prevention of post-thoracotomy pain. *Pain, 48*(3), 299-305.
- Liu, N., Wajima, Z., & Shiga, T. (1999). Opioid-induced bowel dysfunction. *Journal of Clinical Anesthesia, 11*(4), 312-320.
- Perkins, F. M., & Kehlet, H. (2000). Chronic pain as an outcome of surgery. A review. *Anesthesiology, 93*(4), 1123-1133.

Ready, L. B., & Edwards, W. T. (1993). *Management of acute pain: A practical guide*. Mosby Year Book.

Wheeler, J. G., & Smith, C. R. (2002). Opioid-induced respiratory depression in surgical patients. *Anesthesiology Clinics of North America*, 20(4), 843-858.