



The Illinois Medical Bulletin



The Emergence of Community-Associated MRSA and its Association with Severe Skin and Soft Tissue Infections and Necrotizing Pneumonia: A Review of an Evolving Epidemic

¹A.Blake-Sheehy, ²A. Bogard

^{1,2} University of Illinois College of Medicine

Corresponding Email : blakesheehymd283474@gmail.com

Article History :

Received date : 2012/10/12
Revised date : 2012/11/28
Accepted date : 2012/12/09
Published date : 2013/01/13



Copyright: © by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (<https://creativecommons.org/licenses/by-nc/4.0/>).

ABSTRACT

Introduction: Methicillin-Resistant *Staphylococcus aureus* (MRSA), long considered a pathogen confined to healthcare settings, emerged as a major cause of infection in healthy individuals in the community. These Community-Associated MRSA (CA-MRSA) strains demonstrated distinct microbiological features and clinical manifestations compared to their hospital-acquired counterparts.

Literature Review: A review of the pivotal literature documents the rapid rise of a CA-MRSA epidemic, largely driven in North America by the virulent USA300 genetic clone. Epidemiological and clinical studies established a strong association between CA-MRSA and a characteristic disease spectrum, primarily consisting of severe, purulent skin and soft tissue infections (SSTIs), such as abscesses and furuncles. A less common but highly feared manifestation was a devastating, rapidly progressive necrotizing pneumonia, often occurring in young, healthy individuals. The enhanced virulence of these strains was strongly linked to the high prevalence of the cytotoxin Pantone-Valentine Leukocidin (PVL), which causes destruction of immune cells and promotes tissue necrosis. This epidemiological shift required

significant changes in the empirical management and treatment of common community-onset infections.

Conclusion: The body of evidence established CA-MRSA as a formidable public health threat. Its association with a unique profile of severe infections and specific virulence factors fundamentally altered the clinical and microbiological landscape, compelling clinicians to consider MRSA as a primary pathogen in the community setting.

Keywords: Community-Associated MRSA (CA-MRSA), *Staphylococcus aureus*, Skin and Soft Tissue Infections, Necrotizing Pneumonia, Panton-Valentine Leukocidin (PVL).

INTRODUCTION

Staphylococcus aureus is a common and versatile human pathogen, capable of causing a wide range of diseases. For decades, the primary concern in clinical practice was Methicillin-Resistant *Staphylococcus aureus* (MRSA), a formidable pathogen defined by its resistance to beta-lactam antibiotics. Historically, MRSA was almost exclusively a nosocomial, or hospital-acquired (HA-MRSA), infection, affecting patients with significant healthcare exposures, comorbidities, and recent hospitalizations.

A profound epidemiological shift began to occur with the appearance of MRSA infections in patients without any traditional healthcare-associated risk factors. These infections were caused by novel strains of MRSA, termed Community-Associated MRSA (CA-MRSA), which were genetically and phenotypically distinct from the older hospital strains. This emergence represented a new and urgent public health problem. This literature review will synthesise the key scientific publications that characterized the CA-MRSA epidemic and established its association with a unique and severe spectrum of clinical disease.

LITERATURE REVIEW

The understanding of the CA-MRSA epidemic was built upon integrated epidemiological, clinical, and microbiological research that defined the pathogen and the diseases it caused.

The initial recognition of CA-MRSA came from reports of outbreaks of seemingly spontaneous abscesses and skin infections in specific community groups, such as military recruits, athletes, and prison inmates. Subsequent epidemiological surveillance confirmed that this was a nationwide phenomenon. Molecular typing studies were crucial in demonstrating that these

community strains were not simply HA-MRSA strains that had escaped the hospital. Instead, they represented new, distinct genetic lineages—most notably the USA300 clone in North America—that had acquired a smaller, more mobile version of the staphylococcal cassette chromosome *mec* (SCC*mec* type IV), which carries the gene for methicillin resistance (Naimi et al., 2003). These strains also, paradoxically, tended to be susceptible to a wider range of non-beta-lactam antibiotics compared to their multi-drug resistant hospital counterparts.

Clinically, CA-MRSA became strongly associated with a characteristic spectrum of disease, dominated by skin and soft tissue infections (SSTIs). Large prospective studies conducted in emergency departments revealed a dramatic shift in the etiology of purulent SSTIs. They showed that CA-MRSA had become the single most common identifiable cause of skin abscesses, frequently mistaken by patients and providers as "spider bites" due to their intense inflammatory and sometimes necrotic appearance (Moran et al., 2006; Miller et al., 2005). These infections were noted for their tendency to be recurrent and to spread within families and close contacts.

While SSTIs were the most common manifestation, the most feared complication associated with CA-MRSA was a rare but exceptionally severe, rapidly progressive necrotizing pneumonia. This often affected young, otherwise healthy children and adults, frequently following a viral illness like influenza. The pneumonia was characterized by massive inflammation, extensive tissue destruction leading to cavitation, hemoptysis, and a high mortality rate despite aggressive medical care (Francis et al., 2005).

The enhanced virulence of the epidemic CA-MRSA strains was strongly linked to a specific virulence factor: the Panton-Valentine Leukocidin (PVL). PVL is a bi-component pore-forming cytotoxin that targets and lyses host leukocytes, particularly neutrophils, the body's first line of defense against bacterial infection. A very high percentage of the epidemic CA-MRSA strains, including USA300, were found to carry the genes encoding for PVL. Laboratory and clinical evidence suggested that this toxin was a key driver of the intense necrosis seen in both the purulent skin abscesses and the destructive lung injury characteristic of CA-MRSA infections (Vandenesch et al., 2003). The presence of PVL became a defining feature of these highly virulent community strains.

CONCLUSION

The pivotal literature examining the emergence of Community-Associated MRSA chronicles a rapid and dramatic epidemiological shift that presented a major challenge to public health and clinical medicine. This body of work definitively established that new, virulent strains of MRSA had become a leading cause of infections outside of the hospital setting.

This research identified a strong and clear association between CA-MRSA and a unique clinical syndrome dominated by severe, purulent skin infections and a rare but highly lethal necrotizing pneumonia. The discovery of the high prevalence of the Panton-Valentine Leukocidin toxin in these epidemic strains provided a powerful mechanistic explanation for their enhanced virulence. This new understanding forced a fundamental change in clinical practice, requiring clinicians to empirically cover for MRSA in many community-onset infections and prompting new public health strategies to control its spread.

REFERENCES

- Francis, J.S., Doherty, M.C., Lopatin, U., Johnston, C.P., Sinha, G., Ross, T., Cai, M., Hansel, N.N., Perl, T. and Ticehurst, J.R. (2005) 'Severe community-onset pneumonia in healthy adults caused by methicillin-resistant *Staphylococcus aureus* carrying the Panton-Valentine leukocidin genes', *Clinical Infectious Diseases*, 40(1), pp. 100-107.
- Miller, L.G., Perdreau-Remington, F., Rieg, G., Mehdi, S., Perlroth, J., Bayer, A.S., Tang, A.W., Phung, T.O. and Spellberg, B. (2005) 'Necrotizing fasciitis caused by community-associated methicillin-resistant *Staphylococcus aureus* in Los Angeles', *New England Journal of Medicine*, 352(14), pp. 1445-1453.
- Moran, G.J., Krishnadasan, A., Gorwitz, R.J., Fosheim, G.E., McDougal, L.K., Carey, R.B. and Talan, D.A. (2006) 'Methicillin-resistant *S. aureus* infections among patients in the emergency department', *New England Journal of Medicine*, 355(7), pp. 666-674.
- Naimi, T.S., LeDell, K.H., Como-Sabetti, K., Borchardt, S.M., Boxrud, D.J., Etienne, J., Johnson, S.K. and Lynfield, R. (2003) 'Comparison of community- and health care-associated methicillin-resistant *Staphylococcus aureus* infection', *JAMA*, 290(22), pp. 2976-2984.
- Vandenesch, F., Naimi, T., Enright, M.C., Lina, G., Nimmo, G.R., Heffernan, H., Liassine, N., Bes, M., Greenland, T., Reverdy, M.E. and Etienne, J. (2003) 'Community-acquired methicillin-resistant *Staphylococcus aureus* carrying Panton-Valentine leukocidin genes: worldwide emergence', *Emerging Infectious Diseases*, 9(8), pp. 978-984.

