



The Illinois Medical Bulletin



The Hygiene Hypothesis and the Rise of Immunological Diseases: A Review of Epidemiological and Mechanistic Evidence


¹A. Bass, ²A. Beard

^{1,2} University of Illinois College of Medicine

Corresponding Email : bassmd39485@gmail.com

Article History :

Received date : 2009/10/12
Revised date : 2009/11/28
Accepted date : 2009/12/09
Published date : 2010/01/13



Copyright: © by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (<https://creativecommons.org/licenses/by-nc/4.0/>).

ABSTRACT

Introduction: A striking increase in the prevalence of allergic and autoimmune diseases has occurred in industrialized nations over recent decades. The "Hygiene Hypothesis" proposes that this rise is an unintended consequence of reduced exposure to microorganisms in early life, leading to improper calibration of the immune system.

Literature Review: A review of pivotal literature provides strong support for this hypothesis. Large epidemiological studies consistently demonstrated that early-life exposure to microbially rich environments, such as traditional farms, was strongly protective against the development of asthma and atopic sensitization. The mechanistic basis for this observation was extensively explored, leading to a refinement of the original hypothesis. The focus shifted from a simple lack of infections driving a T-helper 1/T-helper 2 cell imbalance to a more nuanced model emphasizing the role of microbial diversity in inducing a robust population of regulatory T-cells (Tregs). These Tregs are essential for establishing and maintaining immune tolerance. Research also identified the critical role of the gut microbiome and exposure to specific microbial products, like endotoxin, in educating the developing immune system.

Conclusion: The body of evidence reviewed confirms a strong association between a sanitized modern lifestyle and the risk of developing immunological disorders. It establishes that the developing immune system requires diverse microbial inputs to mature properly and to establish self-tolerance, and that a lack of such exposure contributes to the rising prevalence of allergies and autoimmunity.

Keywords: Hygiene Hypothesis, Allergy, Autoimmunity, Immune Regulation, Regulatory T-cells (Tregs), Microbiome.

INTRODUCTION

The public health landscape of the last century has been marked by a profound epidemiological transition. While infectious diseases have been progressively controlled in many parts of the world through vaccination, antibiotics, and improved sanitation, a new epidemic has emerged: a dramatic and sustained increase in the prevalence of immune-mediated diseases, including allergic conditions like asthma, hay fever, and eczema, as well as autoimmune disorders such as type 1 diabetes and multiple sclerosis.

To explain this paradoxical inverse relationship, the "Hygiene Hypothesis" was proposed. In its simplest form, this hypothesis posits that a lower incidence of infection in early childhood, resulting from more hygienic living conditions and smaller family sizes, could predispose individuals to the development of immunological diseases later in life. This concept has since evolved significantly, sparking intense research into the complex relationship between the microbial environment and immune system development. This literature review will synthesise the pivotal epidemiological and immunological evidence that explored and refined the Hygiene Hypothesis.

LITERATURE REVIEW

The evidence supporting the Hygiene Hypothesis is drawn from robust epidemiological observations and a progressively more sophisticated understanding of the underlying immunological mechanisms.

The epidemiological support is compelling and consistent. A number of large, cross-sectional, and prospective birth cohort studies provided strong evidence for the protective effect of certain lifestyles and environments. Landmark studies repeatedly found that children raised on traditional farms, with regular exposure to livestock, barns, and unpasteurized milk, had a

remarkably lower prevalence of asthma, allergic rhinitis, and atopic sensitization compared to children living in non-farming rural or urban areas. This "farm effect" was found to be strongly associated with exposure to a higher burden and diversity of environmental microbes, particularly bacterial endotoxin (lipopolysaccharide) found in farm dust (Riedler et al., 2001). Other epidemiological observations, such as the protective effect of having older siblings or attending daycare early in life, further supported the concept that increased microbial exchange in childhood is protective against atopy.

These observations prompted a refinement of the original hypothesis. It became clear that the issue was not simply a lack of specific pathogenic infections, but rather a more general "microbial deprivation" (Yazdanbakhsh et al., 2002). The focus shifted to the idea that the immune system has evolved to expect and require continuous signals from a diverse array of environmental microbes—most of them non-pathogenic—to develop properly. The modern, sanitized lifestyle, with its processed food, purified water, and limited contact with the natural environment, deprives the immune system of these essential developmental cues.

The mechanistic explanation for how microbial exposure educates the immune system also evolved significantly. The initial model proposed a simple balance between T-helper 1 (Th1) cells, which are geared towards fighting intracellular pathogens, and T-helper 2 (Th2) cells, which drive allergic responses. The theory was that insufficient Th1-driving infections led to a default, unopposed dominance of the Th2 pathway. While this model was partially correct, a more critical mechanism was subsequently identified involving a specialized subset of lymphocytes known as regulatory T-cells (Tregs). A large body of research demonstrated that exposure to microbial antigens is a primary stimulus for the induction and expansion of Tregs. These cells are master regulators of the immune system, actively suppressing excessive inflammatory responses from both Th1 and Th2 cells and playing a crucial role in maintaining tolerance to self-antigens and harmless environmental antigens (Bach, 2002).

Further research highlighted the gut as a primary site for this immune education. The trillions of commensal bacteria comprising the gut microbiome were found to be essential for the proper development of gut-associated lymphoid tissue and the generation of a robust Treg population. Disruption of this microbiome in early life was therefore implicated as a key factor in the development of immune dysregulation (Okada et al., 2009).

CONCLUSION

The body of scientific literature exploring the Hygiene Hypothesis provides a compelling explanation for the modern epidemic of allergic and autoimmune diseases. Strong and consistent epidemiological data link a microbially-deprived lifestyle to an increased risk of these conditions. The mechanistic research has moved beyond a simple theory of infection to a sophisticated model of immune regulation, establishing that the human immune system co-evolved with microbes and requires their input for proper calibration.

The evidence confirms that exposure to a diverse range of microorganisms in early life is critical for inducing regulatory T-cell populations that are essential for maintaining immune tolerance. A lack of this "immune education" leaves the system unbalanced and prone to inappropriate inflammatory responses against harmless allergens or self-tissues. This understanding fundamentally reframes immunological diseases, not as a result of a faulty immune system, but as a normal immune system responding predictably to an abnormal, overly hygienic modern environment.

REFERENCES

- Bach, J.F. (2002) 'The effect of infections on susceptibility to autoimmune and allergic diseases', *New England Journal of Medicine*, 347(12), pp. 911-920.
- Liu, A.H. (2007) 'Endotoxin exposure in allergy and asthma: reconciling a paradox', *The Journal of Allergy and Clinical Immunology*, 119(2), pp. 322-329.
- Okada, H., Kuhn, C., Feillet, H. and Bach, J.F. (2009) 'The 'hygiene hypothesis' for autoimmune and allergic diseases: an update', *Clinical & Experimental Immunology*, 160(1), pp. 1-9.
- Riedler, J., Braun-Fahrlander, C., Eder, W., Schreuer, M., Waser, M., Maisch, S., Carr, D., Schierl, R., Nowak, D. and von Mutius, E. (2001) 'Exposure to farming in early life and development of asthma and allergy: a cross-sectional survey', *The Lancet*, 358(9288), pp. 1129-1133.
- Yazdanbakhsh, M., Kremsner, P.G. and van Ree, R. (2002) 'Allergy, parasites, and the hygiene hypothesis', *Science*, 296(5567), pp. 490-494.