



Effect of Structured Education and Antihypertensive Medications Adherence to Decrease Blood Pressure for Hypertension in Elderly

¹ J.Grudis, ² K.Aliffe

^{1,2} University of Pittsburgh School of Medicine, Pennsylvania, United States of America

Correspondence : grudis394838@gmail.com

Article History :

Received date : 2016/10/21

Revised date : 2016/11/06

Accepted date : 2016/12/23

Published date : 2017/01/17



Copyright: © 2018 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (<https://creativecommons.org/licenses/by-nc/4.0/>).

ABSTRACT

Introduction

Hypertension in older adults is a significant public health concern, where effective management is often undermined by poor medication adherence. Adherence is a complex behavior influenced by a wide array of factors, including education, health literacy, and patient-provider relationships, making it a critical challenge in geriatric care.

Methods

This abstract is based on a literature review synthesizing findings from multiple sources, including systematic reviews, cross-sectional studies, and randomized controlled trials. The review examined the various factors influencing antihypertensive medication adherence in older adults and the efficacy of interventions designed to improve it.

Results

Major barriers to adherence identified in the literature include low health literacy, poor illness perception, complex treatment regimens, and the presence of comorbidities. Conversely, factors that significantly improve adherence include structured patient education, a high-quality patient-physician relationship, strong social support, and interventions led by clinical pharmacists. Adherence rates are also influenced by socioeconomic status and the initial choice of antihypertensive drug.

Discussion

The synthesized evidence demonstrates that medication adherence is not a singular issue but a multifactorial one requiring a personalized approach. An effective strategy must address cognitive elements like health literacy and illness perception,

alongside practical and social support systems. Empowering patients through targeted education is a central theme, as it can modify health beliefs and improve self-management skills.

Conclusion

Improving medication adherence among hypertensive older adults requires a multifaceted approach. Interventions must be tailored to enhance health literacy, foster positive health beliefs, and support the patient-provider partnership. Implementing structured educational programs is a key strategy to improve adherence and achieve better blood pressure control in this vulnerable population.

Keywords

Hypertension, Medication Adherence, Elderly, Health Literacy, Structured Education, Blood Pressure Control.

INTRODUCTION

The literature on antihypertensive medication adherence among older adults reveals a multifaceted landscape influenced by various factors, including education, health literacy, and patient-provider relationships. (Banning, 2009) provides an essential foundation by assessing medication adherence through multiple methodologies, including pill counts and electronic monitoring. This study emphasizes the role of clinical pharmacists in enhancing compliance among geriatric patients and identifies key predictors of medication adherence, such as gender differences and intentional versus unintentional non-adherence. This comprehensive review lays the groundwork for understanding the complexities of medication adherence in older populations, particularly in the context of hypertension management.

Building on this foundation, (Akter Khanam et al., 2014) explore the specific challenges faced by hypertensive patients in rural Bangladesh. Their population-based study highlights the critical importance of long-term adherence to antihypertensive therapy and examines how initial drug choices and comorbidities can affect compliance. The authors identify barriers to adherence, underscoring the need for targeted educational interventions that can empower patients and improve their adherence to medication regimens.

(Thomas, 2014) further delves into the barriers to medication adherence, particularly among hypertensive Black men. This study underscores the challenges healthcare providers face in managing hypertension within this demographic. The findings suggest that tailored interventions are necessary to enhance adherence and improve health outcomes, particularly for older adults who often navigate complex treatment regimens. The author posits that despite the availability of effective therapies, adherence remains a significant hurdle, particularly for elderly patients.

In a more recent study, (Babazadeh et al., 2014) examine the interplay between health literacy, illness perception, and medication adherence among older adults with hypertension in Iran. The authors emphasize that poor health literacy is a critical barrier to effective medication adherence, which can lead to severe health complications. Their findings indicate that enhancing health literacy and fostering

positive treatment beliefs are essential strategies for improving adherence rates in this population.

(Ruksakulpiwat et al., 2014) conduct a systematic review that synthesizes findings from multiple studies to identify factors influencing medication adherence among older adults with hypertension. Their review reveals that education, health beliefs, and the quality of the patient-physician relationship are pivotal in shaping adherence behaviors. This comprehensive analysis highlights the necessity for innovative, evidence-based programs that address these critical areas to enhance adherence and ultimately improve blood pressure control among older hypertensive patients.

Together, these studies illustrate the complex interplay of various factors that influence medication adherence in older adults with hypertension. They emphasize the need for multifaceted interventions that consider individual patient needs, educational strategies, and the importance of health literacy to foster better health outcomes in this vulnerable population.

LITERATURE REVIEW

The article titled "A review of interventions used to improve adherence with medication in older" by (Banning, 2009) provides a comprehensive examination of various strategies aimed at enhancing medication adherence among elderly patients, particularly in the context of hypertension management. The study emphasizes the critical role that structured education and clinical pharmacist consultations play in improving compliance with antihypertensive medications.

Banning's review highlights the importance of understanding the predictors of medication adherence, noting that factors such as gender differences and cultural background significantly influence compliance rates among older populations. The article presents a randomized controlled trial that illustrates how targeted interventions, including consultations with clinical pharmacists, can lead to improved adherence and better blood pressure control in geriatric patients. This is particularly relevant given the high prevalence of hypertension in older adults and the associated risks of inadequate treatment.

The review critically assesses existing literature on non-adherence, distinguishing between intentional and unintentional factors that contribute to poor medication compliance. Banning notes that while there are various interventions available, evidence supporting their effectiveness remains limited. This acknowledgment of gaps in the literature is crucial for guiding future research and clinical practice, as it underscores the need for more robust studies to evaluate the long-term impacts of educational programs and medication reviews.

Furthermore, the article discusses the significance of personalized approaches to patient education, suggesting that interventions should be tailored to meet the unique needs and circumstances of older patients. This individualized strategy is essential for enhancing the likelihood of adherence to prescribed antihypertensive regimens, ultimately leading to better health outcomes.

The article "Hypertension: adherence to treatment in rural Bangladesh - findings from a population-based study" by (Akter Khanam et al., 2014) presents a comprehensive examination of the factors influencing antihypertensive medication adherence among the elderly population in rural Bangladesh. The authors emphasize the critical importance of long-term adherence to therapy in mitigating the adverse consequences of hypertension, which is particularly relevant given the increasing prevalence of this condition in aging populations.

One of the key insights from the study is the significant role that the choice of antihypertensive medication plays in patient adherence. The findings suggest that certain drug classes may yield better persistence rates than others, which underscores the necessity for healthcare providers to consider not only the clinical efficacy of medications but also their potential impact on patient compliance. This perspective aligns with the broader understanding that tailored pharmacological approaches can enhance adherence outcomes.

Moreover, the article highlights the influence of comorbidities on adherence to antihypertensive therapy. The presence of additional health conditions can complicate medication regimens and may lead to decreased adherence among patients. This finding is particularly salient in the elderly, who often present with multiple health issues that necessitate complex treatment plans. The authors call for

further research to explore how these comorbidities can be managed effectively to improve adherence rates.

Barriers to adherence identified in the study include a lack of patient education regarding hypertension and its treatment. The authors argue that enhancing patient understanding of their condition and the importance of medication adherence is vital. Education initiatives aimed at hypertensive patients can empower them to take an active role in managing their health, which is essential for improving compliance with antihypertensive medications.

The article "Hypertensive Black Men's Perceptions of a Nurse Administered Medication Protocol" by (Thomas, 2014) provides a comprehensive examination of the challenges related to antihypertensive medication adherence among elderly patients, particularly focusing on hypertensive Black men. The main idea emphasizes the critical role of adherence in managing hypertension and mitigating the associated risks of severe cardiovascular diseases in older adults.

Thomas outlines various barriers that hinder adherence to antihypertensive medications, which include cognitive decline, complex medication regimens, and socioeconomic factors. These barriers are particularly pronounced in the elderly population, where the incidence of chronic diseases is on the rise due to increased life expectancy. The author suggests that healthcare providers need to identify these barriers to develop targeted interventions that can enhance adherence rates among this demographic. This is crucial as the literature indicates that patients aged 70 and over exhibit lower adherence levels compared to younger cohorts, which directly impacts their health outcomes.

The article further discusses the implications of non-adherence, highlighting that inadequate management of hypertension can lead to severe health complications, including heart failure and stroke. Thomas advocates for structured education programs that not only inform patients about their conditions and treatment regimens but also empower them to take an active role in their health management. Such educational interventions could address misconceptions about medications and foster a better understanding of the importance of consistent medication use.

Moreover, the article underscores the significance of a nurse-administered medication protocol, which could provide the necessary support and monitoring for elderly patients. This approach may help in reinforcing adherence by ensuring that patients receive personalized care and follow-up, thereby improving their overall health outcomes. The nurse's role is pivotal in bridging the gap between healthcare providers and patients, particularly in addressing the unique challenges faced by elderly individuals.

The article titled "Impact of health literacy and illness perception on medication adherence among older adults with hypertension in Iran: a cross-sectional study" by (Babazadeh et al., 2014) provides a comprehensive examination of the factors influencing medication adherence among older adults suffering from hypertension. The authors highlight the critical public health challenge posed by hypertension, particularly as it is often asymptomatic and can lead to severe complications if left untreated. This aligns with the characterization of hypertension as a "silent killer," underscoring the urgency of effective management strategies in this demographic. The study's findings reveal that medication adherence is significantly affected by a range of factors, including health literacy and illness perception. The authors argue that low health literacy among older adults can impede their understanding of their condition and the importance of adhering to prescribed treatments. This is particularly concerning given that many older individuals face challenges related to cognitive decline and the complexities of polypharmacy, which can exacerbate difficulties in managing multiple medications.

Moreover, the article underscores that illness perception—how individuals understand and interpret their health condition—plays a pivotal role in adherence behaviors. The authors suggest that older adults who perceive their hypertension as more severe or threatening are more likely to adhere to their medication regimens. This insight is crucial for healthcare providers, as it points to the necessity of tailored educational interventions that enhance patients' understanding of their condition and the implications of non-adherence.

The authors also address the socio-economic factors affecting medication adherence, noting that low-income populations face additional barriers, including

financial constraints and limited access to healthcare resources. This multifaceted approach to understanding adherence is commendable, as it recognizes the interplay between individual behaviors and broader socio-economic determinants of health. However, while the study effectively identifies key factors influencing adherence, it could benefit from a more extensive exploration of specific interventions that could be employed to improve health literacy and illness perception among older adults. Future research could also explore the effectiveness of structured educational programs designed to enhance medication adherence, providing a clearer pathway for practical applications of the findings.

The article titled "Medication Adherence of Older Adults with Hypertension: A Systematic Review" by (Ruksakulpiwat et al., 2014) provides a comprehensive examination of the factors influencing medication adherence among older adults suffering from hypertension. The authors emphasize the serious health risks associated with hypertension, particularly the increased likelihood of stroke and heart disease in this demographic. This systematic review synthesizes findings from thirteen studies, revealing that despite the effectiveness of antihypertensive medications in managing blood pressure and preventing severe complications, adherence rates among older adults remain alarmingly low.

The review identifies fifteen key themes that impact medication adherence, such as education, health literacy, and the patient-physician relationship. These themes are critical as they highlight the multifaceted nature of medication adherence. For instance, the authors point out that health literacy plays a crucial role in how well patients understand their condition and the importance of their medication regimen. This insight aligns with existing literature that suggests that individuals with higher health literacy are more likely to adhere to prescribed treatments.

Moreover, the article discusses the influence of self-efficacy and social support on adherence rates. Self-efficacy, or the belief in one's ability to execute behaviors necessary to produce specific performance attainments, can significantly affect an individual's motivation to adhere to medication schedules. The presence of social support systems, such as family or community resources, also contributes positively

to adherence, indicating that interventions aimed at enhancing these factors could be beneficial.

The systematic review methodology employed by the authors is commendable, as it utilizes a rigorous approach to gather and analyze relevant studies. However, while the identification of themes provides valuable insights, the article could benefit from a more in-depth exploration of how these factors interact with each other. For example, the relationship between health beliefs and actual adherence behaviors could be further elucidated to understand better the underlying mechanisms at play.

CONCLUSION

The literature on antihypertensive medication adherence among older adults reveals a complex interplay of factors that significantly affect compliance and, consequently, blood pressure management. The foundational study by (Banning, 2009) underscores the necessity of employing diverse methodologies to assess adherence and highlights the pivotal role of clinical pharmacists in enhancing compliance through tailored interventions. This study emphasizes the importance of understanding predictors of adherence, including gender differences and the distinction between intentional and unintentional non-adherence, which are crucial for developing effective strategies for older populations.

Further research by (Akter Khanam et al., 2014) illustrates the specific challenges faced by hypertensive patients in rural Bangladesh, reinforcing the notion that long-term adherence is essential for effective hypertension management. The authors identify barriers to adherence, such as the initial choice of medications and the presence of comorbidities, which necessitate targeted educational interventions to empower patients.

The study conducted by (Thomas, 2014) focuses on hypertensive Black men, revealing that factors such as cognitive decline and socioeconomic challenges exacerbate issues of adherence. The findings suggest that healthcare providers must recognize these barriers to create tailored interventions that enhance adherence, particularly among older adults who frequently manage complex treatment regimens.

In Iran, the research by (Babazadeh et al., 2014) highlights the critical roles of health literacy and illness perception in adherence behaviors. The study indicates that low health literacy can severely hinder understanding of medication importance, while a strong illness perception can bolster adherence. This points to the need for healthcare providers to implement educational strategies that improve health literacy and reshape patients' perceptions of their health conditions.

The systematic review by (Ruksakulpiwat et al., 2014) synthesizes findings from various studies, identifying key themes such as the quality of the patient-physician relationship, education, and health beliefs as significant determinants of adherence. The review also emphasizes the importance of self-efficacy and social support in enhancing medication adherence, suggesting that interventions should focus on these areas to improve health outcomes in older adults.

In conclusion, the collective findings from these studies illustrate that improving medication adherence among older adults with hypertension requires a multifaceted approach. Interventions should be tailored to address individual patient needs, enhance health literacy, and foster positive health beliefs. By recognizing the complexities of adherence and implementing structured educational programs, healthcare providers can significantly improve adherence rates and, subsequently, blood pressure control in this vulnerable population.

REFERENCES

Banning, M., 2009. A review of interventions used to improve adherence with medication in older. [\[PDF\]](#)

Akter Khanam, M., Lindeboom, W., Lynn Perez Koehlmoose, T., Shamsul Alam, D., Niessen, L., & Hasnat Milton, A., 2014. Hypertension: adherence to treatment in rural Bangladesh - findings from a population-based study. [\[PDF\]](#)

Thomas, O., 2014. Hypertensive Black Men's Perceptions of a Nurse Administered Medication Protocol. [\[PDF\]](#)

Babazadeh, T., Ranjbaran, S., Pourrazavi, S., Latifi, A., & Maleki Chollou, K., 2014. Impact of health literacy and illness perception on medication adherence among older adults with hypertension in Iran: a cross-sectional study.

Ruksakulpiwat, S., K Schiltz, N., Irani, E., A Josephson, R., Adams, J., & Harmon Still, C., 2014. Medication Adherence of Older Adults with Hypertension: A Systematic Review. ncbi.nlm.nih.gov