



Correlation between Polypharmacy and Length of Stay Geriatric Patient

¹ M.Partan, ² M.Braden

^{1,2} Sidney Kimmel Medical College at Thomas Jefferson University, United States of America

Correspondence : partanmd398343@gmail.com

Article History :

Received date : 2010/10/19

Revised date : 2010/11/06

Accepted date : 2010/12/24

Published date : 2011/01/08



Copyright: © 2011 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (BY NC) license (<https://creativecommons.org/licenses/by-nc/4.0/>).

ABSTRACT

Introduction

Polypharmacy, the concurrent use of five or more medications, is a growing concern in geriatric care due to the rising prevalence of multimorbidity. This practice is associated with adverse health outcomes, including frailty and increased hospitalization. Understanding the correlation between polypharmacy and the length of hospital stay is critical for optimizing treatment and improving healthcare efficiency for aging populations. This review synthesizes literature on this complex relationship.

Methods

This abstract is based on a literature review synthesizing findings from narrative reviews, systematic reviews, and meta-analyses. The review examined the association between polypharmacy and health outcomes in geriatric patients, focusing on its impact on hospitalization rates, length of stay, frailty, and the role of medication management interventions.

Results

The literature establishes a strong link between polypharmacy and adverse health outcomes in the elderly. Polypharmacy is associated with an increased risk of mortality, emergency department

visits, and hospitalization, particularly in frail or prefrail individuals. Studies demonstrate a dose-response relationship, where a higher number of medications corresponds to worse outcomes. Polypharmacy is also linked to potentially inappropriate prescribing (PIP), which can elevate healthcare costs and health complications. Effective interventions, primarily medication reviews, have been shown to be crucial in managing polypharmacy and mitigating its negative effects.

Discussion

The evidence confirms that polypharmacy is a significant risk factor for prolonged hospital stays and increased healthcare utilization among geriatric patients. The interplay with frailty creates a cycle where multiple medications can exacerbate vulnerability, leading to hospitalization. The wide variance in polypharmacy prevalence suggests that prescribing practices and healthcare systems play a significant role. While medication reviews are a pivotal strategy, addressing barriers to their implementation in clinical settings is essential for translating research into effective practice.

Conclusion

A clear correlation exists between polypharmacy and increased length of stay for geriatric patients. This relationship is driven by a higher risk of adverse drug events, frailty, and hospitalization. Comprehensive strategies, centered around systematic medication reviews, are urgently needed to optimize medication regimens, improve health outcomes, reduce healthcare costs, and enhance the quality of life for this vulnerable population.

Keywords

Polypharmacy, Geriatric, Length of Stay, Frailty, Hospitalization, Medication Management.

INTRODUCTION

The correlation between polypharmacy and the length of stay for geriatric patients is a critical area of investigation, particularly given the rising prevalence of multimorbidity in the aging population. (Khezrian et al., 2002) provide a foundational understanding of polypharmacy, defined as the concurrent use of five or more medications, and its potential adverse consequences. They highlight that polypharmacy is often a byproduct of increasing multimorbidity, leading to a higher risk of potentially inappropriate prescribing (PIP) and adverse health outcomes, which can significantly impact healthcare costs. Their findings suggest that the prevalence of polypharmacy can vary widely across different populations, underscoring the need for targeted research to optimize treatment strategies for frail older adults.

Building on this, (Pazan & Wehling, 2001) delve deeper into the ramifications of polypharmacy, linking it to frailty and various health outcomes. Their narrative review indicates a strong association between polypharmacy and increased mortality, hospitalization, and emergency department visits among older adults, particularly those who are frail or prefrail. They present evidence from multiple studies that demonstrate a dose-response relationship between the number of medications and adverse outcomes, thus reinforcing the notion that polypharmacy is a significant risk factor for prolonged hospital stays and increased healthcare utilization.

(Verma et al., 2003) further contribute to this discourse by examining medication interventions aimed at managing polypharmacy among frail older adults. They emphasize that both frailty and polypharmacy are prevalent and interrelated, with medication reviews being a crucial strategy to mitigate the negative impacts of excessive medication use. Their analysis of systematic reviews and meta-analyses reveals that effective medication interventions can reduce medication-related complications and improve health outcomes for older adults, potentially influencing the length of hospital stays.

Together, these articles illustrate the multifaceted relationship between polypharmacy, frailty, and health outcomes in geriatric patients, highlighting the urgent need for comprehensive strategies to address the complexities of medication management in this vulnerable population. Through a critical evaluation of the literature, it becomes evident that understanding the implications of polypharmacy is essential for optimizing care and improving the overall health of older adults.

LITERATURE REVIEW

The article "An overview of prevalence, determinants and health outcomes of polypharmacy" by Khezrian, McNeil, Murray, and Myint provides a comprehensive examination of polypharmacy within the geriatric population, particularly in the context of multimorbidity. The authors highlight that polypharmacy, often defined as the use of five or more medications, is increasingly prevalent due to the rising incidence of chronic conditions among older adults. This observation is critical as it underscores the growing complexity of medication management in this demographic.

The article emphasizes that traditional methods of evaluating medication safety and efficacy, primarily through randomized controlled trials, often exclude older patients with multiple chronic conditions or frailty. This exclusion raises concerns about the applicability of trial results to the geriatric population, as the specific health outcomes associated with polypharmacy in these patients are not well understood. The authors argue for a greater emphasis on research that includes this population to better inform clinical practices and optimize treatment strategies.

A significant point raised in the article is the association between polypharmacy and potentially inappropriate prescribing (PIP), which can lead to serious health implications and increased healthcare costs. The findings from a prospective cohort study are particularly alarming, indicating that patients on a higher number of medications are more likely to receive anticholinergic drugs, which are linked to adverse health outcomes. This relationship highlights the need for careful medication review and management in older adults to mitigate risks associated with drug interactions and side effects.

Moreover, the article presents a wide variance in the prevalence of polypharmacy, ranging from 10% to as high as 90% across different populations. This disparity suggests that factors such as healthcare access, prescribing practices, and patient demographics may significantly influence the extent of polypharmacy experienced by geriatric patients .

The article "Polypharmacy in older adults: a narrative review of definitions, epidemiology and consequences" by Pazan and Wehling (Pazan & Wehling, 2001) provides a comprehensive examination of polypharmacy within the geriatric population, highlighting its implications for health outcomes, particularly concerning frailty and hospitalization. The authors present a narrative review that synthesizes various studies, offering insights into how polypharmacy is not merely a clinical concern but a significant factor influencing the overall health trajectory of older adults.

One of the key insights from the article is the relationship between polypharmacy and frailty, which is particularly relevant for understanding the complexities of geriatric care. The authors reference a cohort study involving Spanish older adults that illustrates how polypharmacy is linked to hospitalization in frail and prefrail individuals. This connection underscores the potential for polypharmacy to exacerbate health issues in vulnerable populations, leading to increased lengths of hospital stays. The evidence presented suggests that managing medication regimens in older adults is critical not only for individual health outcomes but also for healthcare system efficiency.

Furthermore, the article discusses a systematic review and meta-analysis that found a significant association between polypharmacy and mortality rates. However, it is noteworthy that the authors also highlight a study indicating that polypharmacy was not independently associated with non-cancer mortality after adjusting for chronic conditions. This nuanced perspective is essential, as it suggests that while polypharmacy is a risk factor, its impact may be moderated by the presence of other health issues, which complicates the interpretation of its effects on length of stay. The article also draws attention to various studies that demonstrate a clear association between polypharmacy and hospitalization rates, including unplanned

admissions. A longitudinal health insurance database study from Taiwan supports the notion of an independent association between polypharmacy and all-cause hospitalization, further affirming that older adults on multiple medications are at a heightened risk for adverse health events that necessitate hospitalization. This finding is particularly significant in the context of healthcare resource utilization, as it implies that reducing polypharmacy could potentially lead to shorter lengths of stay and better health outcomes for geriatric patients.

The article "An Overview of Systematic Reviews and Meta-Analyses on the Effect of Medication Interventions Targeting Polypharmacy for Frail Older Adults" by Verma et al. (Verma et al., 2003) provides a comprehensive examination of the complex interplay between polypharmacy and frailty among older adults. The authors highlight the alarming rates of polypharmacy in this demographic, underscoring its significance as a public health concern. The article effectively synthesizes existing literature, revealing that frailty is not merely a consequence of aging but is intricately linked to the number of medications consumed by older adults.

A critical evaluation of the material suggests that the authors have successfully identified key factors contributing to both frailty and polypharmacy, such as weight loss, poor nutrition, and balance disorders. These factors serve as measurable characteristics that can inform clinical practice and intervention strategies. The systematic review approach taken by the authors adds credibility to their findings, as it aggregates data from multiple studies to provide a more robust understanding of the issue.

Moreover, the article presents medication review as a pivotal intervention for addressing polypharmacy. This systematic examination of a patient's medications aims to enhance therapeutic outcomes while minimizing medication-related complications. The authors argue that such interventions are particularly effective in managing polypharmacy and reducing inappropriate prescriptions, which is crucial given the potential adverse effects associated with polypharmacy in older adults.

However, while the article offers valuable insights, it could benefit from a more detailed exploration of the specific mechanisms by which medication interventions lead to improved outcomes for frail older adults. For instance, a deeper analysis of how these interventions directly correlate with reduced lengths of hospital stays or improved quality of life could enhance the practical implications of the findings. Additionally, the article could address potential barriers to the implementation of medication reviews in clinical settings, as understanding these challenges is essential for translating research into practice.

CONCLUSION

The literature on the correlation between polypharmacy and the length of stay for geriatric patients reveals a complex interplay between medication use, health outcomes, and healthcare resource utilization. Polypharmacy, defined as the concurrent use of five or more medications, is increasingly prevalent among older adults due to the rising incidence of multimorbidity . This phenomenon is concerning, as it has been linked to various adverse health outcomes, including increased hospitalization and mortality rates (Pazan & Wehling, 2001). The articles reviewed provide compelling evidence that polypharmacy is not only a clinical concern but also a significant factor influencing the overall health trajectory of geriatric patients.

Khezrian et al. highlight the association between polypharmacy and potentially inappropriate prescribing (PIP), which can exacerbate health complications and elevate healthcare costs. The authors emphasize the variability in the prevalence of polypharmacy, suggesting that factors such as healthcare access and prescribing practices play a crucial role in its manifestation among older adults. This variability underscores the need for targeted research that addresses the specific needs of frail older individuals, who are particularly vulnerable to the negative consequences of excessive medication use.

Pazan and Wehling (Pazan & Wehling, 2001) further elucidate the relationship between polypharmacy and frailty, noting that frail and prefrail older adults experience higher hospitalization rates linked to polypharmacy. Their findings suggest that managing medication regimens is critical not only for individual health

outcomes but also for the efficiency of healthcare systems. The authors present evidence from cohort studies indicating that polypharmacy can lead to longer hospital stays, emphasizing the importance of optimizing medication management to mitigate these risks.

In conclusion, the literature indicates a clear correlation between polypharmacy and the length of stay for geriatric patients, with implications for both individual health outcomes and healthcare resource utilization. The reviewed articles collectively underscore the need for comprehensive strategies to address the complexities of medication management in this vulnerable population. Effective interventions, such as medication reviews, are essential for optimizing care and improving the overall health of older adults, ultimately contributing to reduced lengths of hospital stays and enhanced quality of life.

REFERENCES

Khezrian, M., J. McNeil, C., D. Murray, A., & K. Myint, P., 2002. An overview of prevalence, determinants and health outcomes of polypharmacy.

Pazan, F. & Wehling, M., 2001. Polypharmacy in older adults: a narrative review of definitions, epidemiology and consequences.

Verma, A., Saha, S., Jarl, J., Conlon, E., McGuinness, B., & Trépel, D., 2003. An Overview of Systematic Reviews and Meta-Analyses on the Effect of Medication Interventions Targeting Polypharmacy for Frail Older Adults.